


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1963

[Full Table of Content](#)



63/R/AG/A/4 Agdell - Grass

Rothamsted Research

Rothamsted Research (1964) *63/R/AG/A/4 Agdell - Grass* ; Yields Of The Field Experiments 1963, pp 17 - 17 - DOI: <https://doi.org/10.23637/ERADOC-1-183>

63/A/4

GRASS - AGDELL 1963

(AG)

For history, treatments, etc., see 'Details of the Classical and Long Term Experiments' 1956.

The areas carrying strip crops in 1961 and 1962 were bare fallowed.

Area harvested (acres): Plots 1, 2 and 3 - 0.0165, plots 4, 5 and 6 - 0.0023.

Cultivations, etc.:

Grass. 'Nitro-Chalk' applied: Mar 25, 1963. Cut: June 6. Grass destroyed with paraquat at 2 lb in 40 gallons per acre: June 15. Fallow areas. Ploughed: Apr 8, 1963.

Note: The grass was abandoned after the first cut because of winter killing and weeds.

Summary of Results

Grass, Dry matter: cwt per acre

Manure to turnips until 1948 Plot Rotation	Mineral manure* no nitrogen				Mineral* and nitrogenous manure+		Mean
	None since 1848 5 Fallow	6 Clover	3 Fallow	4 Clover	1 Fallow	2 Clover	
	3.8	2.1	4.5	5.6	2.1	6.1	4.1

Mean dry matter % as cut: 25.2

*P, K, Na, Mg.

+Rape dust (or castor meal) + ammonium sulphate.