

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1964

[Full Table of Content](#)



64/W/WIR/B/6 Irrigation

Rothamsted Research

Rothamsted Research (1965) *64/W/WIR/B/6 Irrigation ; Yields Of The Field Experiments 1964*, pp 111 - 117 - DOI: <https://doi.org/10.23637/ERADOC-1-160>

64/B/6.1

IRRIGATION EXPERIMENT

(WIR)

Revised 1963

The effects of irrigation and nitrogen - Woburn Butt Close 1964, the 14th year.

For details of previous cropping, treatments etc., see 'Details' 1962.

Sugar beet: The test of early v normal singling was omitted.

Barley: The variety is now Maris Badger.

Clover: The variety is now Dorset Marl double cut red clover.

Lucerne: Plots 73 - 80 failed and were rotary cultivated in mid-season. In view of the incomplete replication remaining no yields are presented.

Area harvested: Sugar beet, sub plot - 0.0102, barley, whole plot - 0.0097, clover, whole plot - 0.0058, lucerne, sub plot - 0.0162.

RAINFALL AND IRRIGATION: INCHES

Week ending	Rain-fall	Barley C	Sugar beet			Clover			Lucerne		
			A	B	C	A	B	C	A	B	C
May 4	0.77										
11	0.09										
18	0.12	0.50				0.50		0.50			
25	0.44					0.50		0.50	0.50		0.50
June 1	0.25	0.50	0.50		0.50						
8	1.51										
15	0.66										
22	1.28										
29	-										
July 6	-	0.50	0.50		0.50		0.50	0.50		0.50	0.50
13	0.55	0.50	0.50		0.50		0.50	0.50		0.50	0.50
20	0.33				0.50		0.50	0.50			
27	0.09										0.50
Aug 3	0.01				1.50	0.50					
10	0.50										
17	0.12						0.50	0.50			
24	0.16				0.50	1.00		0.50	0.50		1.00
31	0.03						0.50	0.50			
Sept 7	0.10										
14	0.01										
21	0.37										
28	0.62										
Oct 5	-										
Total	8.01	2.00	1.50	2.00	3.50	1.00	3.00	4.00	10.50	1.00	3.00

64/B/6.2

Cultivations, etc.:

Sugar beet. Subsoiled: Sept 11, 1963. Ploughed twice: Aug 20 and Jan 2, 1964. Salt applied: Feb 4. Basal compound and sulphate of ammonia applied: Apr 7. Seed drilled at 5lb: Apr 10. Singled, sprayed with DDT at 0.63 lb in 40 gals: May 21. Lifted: Oct 13.

Barley: Ground chalk applied at 44 cwt: Dec 20, 1963. Ploughed: Jan 3, 1964. Basal compound and 'Nitro-Chalk' applied, seed drilled at 2.5 bushels: Feb 14. Clover sown at 30 lb: Apr 27. Sprayed with MCPB/MCPA (New Legumex at 5 pints in 40 gals): May 21. Combine harvested: Aug 21.

Clover: Seed undersown in barley at 30 lb: Apr 18, 1963. Basal compound fertiliser applied: Feb 5, 1964. Cut 3 times: June 11, July 27, Sept 29.

Lucerne: Sprayed with paraquat at 1.6 lb ion in 40 gals: Feb 10, 1964. 'Nitro-Chalk', muriate of potash and basal P applied: Apr 3. Plots 73 - 80 cut (no yields taken) and rotary cultivated twice: May 15 and 27. Cut 4 times: June 9, July 16, Sept 2, Oct 30. Muriate of potash applied after first 3 cuts.

Standard errors per plot.

Sugar beet. Roots,	Whole plot:	1.421 or 8.6% (6 d.f.)
	Sub plot:	1.592 or 9.7% (8 d.f.)
Total sugar,	Whole plot:	6.33 or 9.8% (6 d.f.)
	Sub plot:	6.39 or 9.9% (8 d.f.)
Tops,	Whole plot:	1.738 or 19.4% (6 d.f.)
	Sub plot:	1.364 or 15.3% (8 d.f.)
Barley. Grain,	Whole plot:	5.15 or 15.3% (5 d.f.)
	Sub plot:	2.57 or 7.6% (10 d.f.)
Clover. Dry matter, 1st cut.	Whole plot:	2.24 or 7.5% (6 d.f.)
	Sub plot:	2.71 or 9.0% (8 d.f.)
2nd cut.	Whole plot:	2.51 or 11.5% (6 d.f.)
	Sub plot:	2.50 or 11.4% (8 d.f.)
3rd cut.	Whole plot:	0.86 or 12.0% (6 d.f.)
	Sub plot:	0.55 or 7.7% (8 d.f.)
Total of 3 cuts.	Whole plot:	4.29 or 7.3% (6 d.f.)
	Sub plot:	4.81 or 8.1% (8 d.f.)

64/B/6.3

SUMMARY OF RESULTS

SUGAR BEET

	O	A	B	C	Mean
ROOTS					
	(1) and (2)				(± 0.460)
N1	13.41	13.34	17.48	16.60	15.20
N2	13.75	16.77	19.91	20.17	17.65
Mean (± 0.820)	13.58	15.05	18.69	18.39	16.43
SUGAR %					
N1	20.1	20.1	19.6	19.9	19.9
N2	18.6	19.3	19.3	19.6	19.2
Mean	19.4	19.7	19.5	19.8	19.6

- (1) (± 0.919) For use in vertical and interaction comparisons only
 (2) (± 1.047) For use in horizontal and diagonal comparisons only

64/B/6.4

SUGAR BEET					
	O	A	B	C	Mean
TOTAL SUGAR					
		(1) and (2)			(±1.84)
N1	54.0	53.5	68.5	66.1	60.5
N2	51.5	64.7	76.9	79.4	68.1
Mean (±3.66)	52.8	59.1	72.7	72.7	64.3

TOPS					
		(1) and (2)			(±0.394)
N1	5.91	7.09	8.05	7.87	7.23
N2	7.22	9.54	11.94	13.91	10.65
Mean (±1.004)	6.56	8.31	10.00	10.89	8.94

Total sugar Tops

- (1) (±3.69) (±0.787) For use in vertical and interaction comparisons only
- (2) (±4.49) (±1.148) For use in horizontal and diagonal comparisons only

64/B/6.5

BARLEY
GRAIN

	O	C	Mean
	(1) and (2)		(±0.74)
N1	28.8	31.0	29.9
N2	37.4	37.6	37.5
Mean (±2.10)	33.1	34.3	33.7

(1) (±1.05) For use in vertical and interaction comparisons only
 (2) (±2.23) For use in horizontal and diagonal comparisons only

Mean D.M.%: 83.5

64/B/6.6

CLOVER
DRY MATTER

	O	A	B	C	Mean
1ST CUT*					
		(1) and (2)			(±0.78)
N1	29.8	30.9	28.6	33.2	30.6
N2	27.3	28.0	27.0	34.9	29.3
Mean (±1.29)	28.5	29.5	27.8	34.0	30.0

Mean D.M. %: 17.1

2ND CUT

		(1) and (2)			(±0.72)
N1	17.1	17.5	28.7	23.6	21.7
N2	18.6	15.1	29.0	25.4	22.0
Mean (±1.45)	17.8	16.3	28.9	24.5	21.9

Mean D.M. %: 18.6

1ST CUT 2ND CUT

- (1) (±1.57) (±1.44) For use in vertical and interaction comparisons only
 (2) (±1.70) (±1.77) For use in horizontal and diagonal comparisons only

*NOTE: 1st cut O = B
 A = C

64/B/6.7

CLOVER					
DRY MATTER					
	O	A	B	C	Mean
3RD CUT					
		(1) and (2)			(±0.16)
N1	2.2	3.2	11.9	11.7	7.2
N2	2.4	4.6	10.3	11.6	7.2
Mean (±0.50)	2.3	3.9	11.1	11.6	7.2

Mean D.M. %: 24.1

TOTAL OF 3 CUTS					
		(1) and (2)			(±1.39)
N1	49.1	51.6	69.2	68.5	59.6
N2	48.3	47.7	66.3	71.9	58.5
Mean (±2.48)	48.7	49.7	67.7	70.2	59.1

Mean D.M. %: 19.9

	3RD CUT	TOTAL OF 3 CUTS	
(1)	(±0.32)	(±2.78)	For use in vertical and interaction comparisons only
(2)	(±0.55)	(±3.16)	For use in horizontal and diagonal comparisons only