

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1964

[Full Table of Content](#)



64/R/NL/B/8 N Levels and Residues Two Course Rotation

Rothamsted Research

Rothamsted Research (1965) *64/R/NL/B/8 N Levels and Residues Two Course Rotation ; Yields Of The Field Experiments 1964*, pp 123 - 124 - DOI: <https://doi.org/10.23637/ERADOC-1-160>

64/B/8.1

N LEVELS AND RESIDUES ROTATION

(NL)

Direct and residual effects of sulphate of ammonia - Long Hoos III, 1964, the fifth and final year.

Rotation: Wheat, potatoes. In 1964, winter wheat only, following potatoes.

Design: 3 x 3 x 3 in 3 blocks of 9 plots each.

Area of each plot: 0.0212. Area harvested: 0.0141.

Treatments: All combinations of:-

N 1964.	None (N0), 0.5 (N1), 1.0 (N2) cwt N.
N to potatoes 1963.	None (NP0), 0.75 (NP1), 1.50 (NP2) cwt N.
N to wheat 1960 & 1962.	None (NW0), 0.5 (NW1), 1.0 (NW2) cwt N.

The nitrogen was applied as sulphate of ammonia. Ground chalk was applied in October 1962 at 1 cwt per cwt of sulphate of ammonia applied in 1960, 1961 and 1962.

Basal dressing: 2.25 cwt 0:14:28 combined drilled.

Cultivations, etc.: Chisel ploughed: Oct 14, 1963. Seed drilled at 3 bushels: Oct 19. Sulphate of ammonia applied, plots sprayed with mecoprop/2,4-D (Methoxone Extra at 7 pints in 40 gals): Apr 27, 1964. Combine harvested: Aug 25. Variety: Cappelle.

NOTES: (1) For details of the previous years' results see 'Results' 60/B/10, 61/B/9, 62/B/9 and 63/B/9.

(2) If the N levels to potatoes 1961 are included, the design becomes a 1/3 replicate, but no account of this factor has been taken in the analysis.

Standard error per plot. Winter wheat:
Grain: 2.79 or 10.4% (15 d.f.)

64/B/8.2

SUMMARY OF RESULTS

GRAIN

		NP			NW			Mean	
		0	1	2	0	1	2		
		(±1.61)			(±1.61)			(±0.93)	
N	0	19.0	20.3	25.5	20.9	21.0	22.9	21.6	
	1	24.5	26.1	25.6	25.7	22.9	27.7	25.4	
	2	29.8	35.3	35.6	35.0	32.3	33.4	33.6	
					0	24.7	23.4	25.2	24.4
					NP 1	25.6	26.8	29.2	27.2
					2	31.2	26.0	29.4	28.9
Mean (±0.93)					27.2	25.4	28.0	26.8	

Mean D.M. %: 82.4