

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1964

[Full Table of Content](#)



64/R/LM/C/1 and 64/W/WAC/C/1 Levels of K and Mg - Potatoes and Sugar Beet

Rothamsted Research

Rothamsted Research (1965) *64/R/LM/C/1 and 64/W/WAC/C/1 Levels of K and Mg - Potatoes and Sugar Beet* ; Yields Of The Field Experiments 1964, pp 137 - 142 - DOI:

<https://doi.org/10.23637/ERADOC-1-160>

64/C/1.1

EFFECT OF K AND Mg

(IM and WAC)

K and Mg - Rothamsted (R) Sawyers I the sixth year, potatoes 1964 and Woburn (W) Stackyard Series C the fifth year, potatoes and sugar beet 1964.

Design: Sawyers I (R): 8 randomised blocks of 9 plots each.
Stackyard Series C (W): 4 randomised blocks of 9 plots each, 2 blocks per crop.

Area of each plot:	Area harvested:
Sawyers I (R): 0.0206	0.0137
Stackyard Series C (W): 0.0011	0.0011

Treatments. All combinations of:-

1. Mg: None (Mg 0), 29 (Mg 1), 58 lb (Mg 2) applied as magnesium sulphate on Sawyers I (R) and none (Mg 0), 29 (Mg 1), 58 lb (Mg 2) as kieserite on Stackyard Series C (W).

2. K: Sawyers I (R): (in lb K).

	K0	K1	K2
1962	24	95	165
1963	None	71	142
1964	None	91	182

Stackyard Series C (W): None (K0), 95 (K1), 190 lb (K2), applied in 1962, 1963 and 1964. All K as sulphate of potash.

The K and Mg dressings on Stackyard Series C (W) were applied half in winter half in spring.

In addition in 1959 and 1962 magnesium-free calcium carbonate was applied to blocks on Sawyers I (R) at 38, 76 cwt (CA1, CA2).

Basal dressings:

Sawyers I (R): 0.5 cwt P2O5 as triple superphosphate applied on flat before planting.

1.0 cwt N as sulphate of ammonia applied on flat before planting.

Stackyard Series C (W): 1.5 cwt P2O5 as triple superphosphate, half dug in in winter, half applied in spring.

1.5 cwt N as ammonium nitrate in seedbed.

Cultivations, etc.:

Sawyers I (R): Ploughed: Jan 9, 1964. Fertilisers applied: May 8.

Rotary cultivated, potatoes machine planted: May 11. Earthed

up: June 24. Sprayed 3 times with mancozeb at 1.2 lb in 35 gals:

64/C/1.2

June 30, July 17, July 31. Sprayed with diquat (Reglone at 3 pints in 40 gals): Sept 18. Lifted: Sept 25. Variety: King Edward.

Stackyard Series C (W):

Potatoes: First dressings of P,K and Mg applied, plots dug: Dec 2, 1963. Second dressings of PK and Mg and all N applied: Apr 23, 1964. Rotary cultivated, potatoes planted: Apr 28. Sprayed with copper oxychloride fungicide at 2.3 lb Cu in 40 gals: July 3. Sprayed with dimethoate at 3 fluid oz in 40 gals: July 6. Sprayed with copper oxychloride fungicide at 2.3 lb Cu in 40 gals: July 17. Lifted: Sept 9. Variety: King Edward.

Sugar beet: First dressing of P,K and Mg applied, plots dug: Dec 2, 1963. Second dressing of P,K and Mg and all N applied, plots rotary cultivated, seed drilled at 7 lb: Apr 23, 1964. Singled: June 4. Sprayed with dimethoate at 3 fluid oz in 40 gals: July 6. Lifted: Oct 13. Variety: Klein E.

NOTE: For details of previous years' results see 'Results' 60/Ci/3, 61/C/7, 62/C/6, 63/C/1.

Standard errors per plot.

Sawyers I (R):

Potatoes, total tubers: 0.724 or 10.9% (48 d.f.)

Stackyard Series C (W):

Potatoes, total tubers: 0.805 or 7.8% (8 d.f.)

Sugar beet, roots: 1.262 or 11.1% (8 d.f.)
total sugar: 4.500 or 11.5% (8 d.f.)
tops: 0.772 or 8.6% (8 d.f.)

64/C/1.3

SUMMARY OF RESULTS

Sawyers I (R)

POTATOES, TOTAL TUBERS

	K0	K1	K2	MG0	MG1	MG2	Mean
	(± 0.209)*			(± 0.209)*			
CA1	3.05	7.97	9.51	6.76	7.07	6.70	6.84
CA2	2.12	7.30	9.84	6.43	6.41	6.42	6.42
Diff.	-0.93	-0.67	+0.33	-0.33	-0.66	-0.28	-0.42
		(± 0.295)**			(± 0.295)**		
					(± 0.256)		(± 0.148)
		K0		2.78	2.73	2.25	2.59
		K1		7.42	7.61	7.87	7.64
		K2		9.58	9.89	9.57	9.68
		Mean (± 0.148)		6.59	6.74	6.56	6.63

* For use in horizontal and interaction comparisons only.

** For use only in testing the differences of 2 differences.

64/C/1.4

Sawyers I (R)

POTATOES, % WARE

	K0	K1	K2	MG0	MG1	MG2	Mean
CA1	67.5	89.0	92.1	82.8	84.0	81.8	82.9
CA2	53.2	89.2	92.9	80.8	78.4	76.2	78.5
Diff.	-14.3	+0.2	+0.8	-2.0	-5.6	-5.6	-4.4

K0	65.6	60.8	54.8	60.4
K1	87.0	90.1	90.2	89.1
K2	92.7	92.8	92.1	92.5
Mean	81.8	81.2	79.0	80.7

64/C/1.5

Stackyard Series C (W)

POTATOES, TOTAL TUBERS

	MGO	MG1	MG2	Mean
		(±0.569)		(±0.328)
K0	4.96	5.01	5.31	5.09
K1	10.67	12.55	12.01	11.74
K2	13.59	14.09	14.68	14.12
Mean (±0.328)	9.74	10.55	10.67	10.32

SUGAR BEET, ROOTS

		(±0.893)		(±0.515)
K0	5.63	7.47	7.15	6.75
K1	11.81	12.35	14.16	12.77
K2	14.69	14.37	14.89	14.65
Mean (±0.515)	10.71	11.40	12.07	11.39

64/C/1.6

Stackyard Series C (W)

SUGAR BEET, SUGAR %

	MG0	MG1	MG2	Mean
K0	16.5	16.5	16.6	16.5
K1	17.3	17.3	17.6	17.4
K2	17.1	17.7	17.4	17.4
Mean	17.0	17.2	17.2	17.1

SUGAR BEET, TOTAL SUGAR

		(±3.18)		(±1.84)
K0	18.5	24.6	23.7	22.3
K1	40.8	42.7	49.8	44.4
K2	50.3	50.9	51.8	51.0
Mean (±1.84)	36.6	39.4	41.8	39.3

SUGAR BEET, TOPS

		(±0.546)		(±0.315)
K0	6.20	7.19	7.94	7.11
K1	9.43	9.03	10.32	9.59
K2	10.52	10.22	10.17	10.30
Mean (±0.315)	8.71	8.81	9.48	9.00