

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1964

[Full Table of Content](#)



64/R/LL/C/3 and 64/W/WLL/C/3 Long-term Liming - Spring Beans

Rothamsted Research

Rothamsted Research (1965) *64/R/LL/C/3 and 64/W/WLL/C/3 Long-term Liming - Spring Beans* ; Yields Of The Field Experiments 1964, pp 145 - 147 - DOI:

<https://doi.org/10.23637/ERADOC-1-160>

64/C/3.1

LONG TERM LIMING EXPERIMENT - SPRING BEANS 1964

(LL and WLL)

For treatments etc., see 'Results' 63/C/3 and for previous years' results see 62/C/8 and 63/C/3.

Area of each sub plot: 0.0145. Area harvested: Sawyers I (R) - 0.0133, Stackyard Series C (W) - 0.0096.

Plots receiving both P and K were subdivided for a comparison between broadcasting of powder fertiliser (B) (rates and materials as before) and placement drilling of 409 lb 0:14:28(D).

Basal dressing: 0.25 cwt N as 'Nitro-Chalk' broadcast.

Cultivations, etc.

Sawyers I (R): Ploughed: Nov 14, 1963. Fertilisers applied: Feb 15, 1964. Seed drilled at 200 lb: Mar 6. Sprayed with simazine at 1 lb in 40 gals: Mar 13. Combine harvested: Aug 25. Variety: Spring Tick.

Stackyard Series C (W): Ploughed: Nov 1, 1963. Fertilisers applied and seed drilled* at 200 lb: Nov 8. Sprayed with simazine at 1 lb in 40 gals: Nov 9. Seed drilled* at 200 lb: Mar 13, 1964. Sprayed with simazine at 1 lb in 40 gals: Apr 6. Combine harvested: Aug 25. Variety: Spring Tick.

* The winter beans failed through bird damage and the spring beans were sown without seedbed cultivation.

NOTE: On Sawyers I (R) samples were taken at harvest for pod and bean counts.

Standard error per plot. Grain:

Main experiment

Sawyers I (R): 3.56 or 20.2% (15 d.f.)

Stackyard Series C (W): 1.62 or 10.5% (15 d.f.)

PK plots

Sawyers I (R): Whole plot: 2.46 or 12.4% (3 d.f.)

Sub plot: 0.96 or 4.8% (4 d.f.)

Stackyard Series C (W): Whole plot: 1.30 or 6.6% (3 d.f.)

Sub plot: 2.23 or 11.4% (4 d.f.)

Ground chalk tons per acre.

CA0 = None

CA2 = 2

CA4 = 4

CA8 = 8

64/C/3.2

SUMMARY OF RESULTS

SAWYERS I (R)

GRAIN

Main experiment

	CA0	CA2	CA4	CA8	Mean
Mean (± 1.26)	14.7	19.0	19.8	17.1	17.7
		(± 1.78)			(± 0.89)
-	15.1	20.2	19.7	16.2	17.8
P	14.4	17.8	19.9	18.0	17.5
-	15.2	16.6	17.5	14.7	16.0
K	14.2	21.4	22.0	19.5	19.3
		- P			
	(± 1.26)				
-	16.7	15.3			
K	18.9	19.7			

Mean D.M. %: 83.0

PK plots

	CA0	CA2	CA4	CA8	Mean
		(1) and (2)			(± 0.34)
B	15.1	21.6	22.8	19.2	19.7
D	15.9	21.5	22.4	20.9	20.1
Mean (± 1.23)	15.5	21.5	22.6	20.0	19.9

Mean D.M. %: 82.5

- (1) (± 0.68) For use in vertical and diagonal comparisons
 (2) (± 1.80) For use in horizontal and interaction comparisons

64/C/3.3

STACKYARD SERIES C (W)

GRAIN

Main experiment

	CA0	CA2	CA4	CA8	Mean
Mean (± 0.57)	19.1	16.5	13.0	13.2	15.5
	(± 0.81)				(± 0.41)
-	18.1	16.3	12.4	12.4	14.8
P	20.0	16.7	13.6	14.1	16.1
-	16.6	12.8	10.9	10.7	12.7
K	21.6	20.3	15.2	15.7	18.2
	-	P			
	(± 0.57)				
-	13.1	12.4			
K	16.5	19.8			

Mean D.M. %: 83.9

PK Plots

	CA0	CA2	CA4	CA8	Mean
	(1) and (2)				(± 0.79)
B	22.1	22.3	17.8	17.2	19.8
D	22.3	21.4	17.4	16.5	19.4
Mean (± 0.65)	22.2	21.8	17.6	16.9	19.6

Mean D.M. %: 83.6

- (1) (± 1.58) For use in vertical and diagonal comparisons
 (2) (± 1.44) For use in horizontal and interaction comparisons