

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1964

[Full Table of Content](#)



### **64/R/AY/C/5 and 64/W/WBC/C/5 Methods of Application of Fertiliser 1963-64 - Potatoes**

#### **Rothamsted Research**

Rothamsted Research (1965) *64/R/AY/C/5 and 64/W/WBC/C/5 Methods of Application of Fertiliser 1963-64 - Potatoes* ; Yields Of The Field Experiments 1964, pp 153 - 155 - **DOI:**

**<https://doi.org/10.23637/ERADOC-1-160>**

64/C/5.1

METHODS OF APPLICATION OF FERTILISER 1964 - 65

(AY and WBC)

Methods of application of fertiliser - Rothamsted (R) Whittlocks and Woburn (W) Broad Mead II, 1964 the first year - potatoes.

Design (each field): 3 randomised blocks of 12 plots each.

Area of each plot: 0.0212. Area harvested: 0.0133.

Treatments: None (L0) (3 plots per block) and all combinations of:-

1. Levels of 13:13:20 to supply (cwt):

N	P2O5	K2O
0.66	0.66	1.02 (L1)
1.32	1.32	2.03 (L2)
2.00	2.00	3.07 (L3)

2. Methods of application: Broadcast (B), placed (P), broadcast and rotary cultivated in (BR).

NOTE: The experiment is designed to include an additional factor applied to the 1965 wheat crop, viz. PK broadcast on seedbed and 'Nitro-Chalk' as spring dressing to supply:-  
N, P2O5 and K2O at levels as (0), (1), (2) above.

Basal dressing: None.

Cultivations, etc.:

Whittlocks (R). Ploughed: Oct 11, 1963. Fertilisers applied to ploughing: Apr 8, 1964. Rotary cultivated, fertilisers applied, potatoes planted: Apr 15. Earthed up: June 12. Sprayed three times with mancozeb at 1.2 lb in 35 gals: June 30, July 16, July 31. Sprayed with diquat (Reglone at 3 pints in 40 gals): Sept 8. Lifted: Sept 15. Variety: King Edward. Previous crops: Winter and spring wheat 1962, spring wheat 1963.

Broad Mead (W). Ploughed twice: Oct 2, 1963 and Jan 13, 1964. Fertilisers applied to ploughing: Apr 9. Rotary cultivated: Apr 13. Fertilisers applied, potatoes planted: Apr 14. Sprayed with diquat/paraquat (Preaglone Extra at 2 pints in 40 gals): May 12. Sprayed with mancozeb at 1.6 lb in 50 gals: June 25, and again at 1.2 lb in 36 gals: July 14. Sprayed with diquat (Reglone at 3 pints in 40 gals): Sept 2. Lifted: Sept 10. Variety: King Edward. Previous crops: Spring wheat 1962, barley 1963.

Standard errors per plot. Total tubers:

Whittlocks (R):	0.610 or 5.6% (24 d.f.)
Broad Mead II (W):	1.985 or 17.4% (24 d.f.)

64/C/5.2

SUMMARY OF RESULTS

WHITTLOCKS (R)

	L0	L1	L2	L3	Mean
TOTAL TUBERS					
		(±0.352)			(±0.203)
B		11.23	12.83	13.92	12.66
P		10.72	13.04	13.27	12.34
BR		10.27	13.01	14.60	12.63
Mean (±0.209)	6.18	10.74	12.96	13.93	10.95*

	% WARE				
B		90.4	90.1	91.4	90.6
P		86.3	91.7	91.4	89.8
BR		88.3	91.9	91.1	90.4
Mean	82.0	88.3	91.2	91.3	88.2*

\* General mean

64/C/5.3

BROAD MEAD II (W)

	L0	L1	L2	L3	Mean
TOTAL TUBERS					
(±1.146)					
B		9.82	11.90	14.84	12.19
P		12.96	15.55	14.09	14.20
BR		10.06	13.50	14.66	12.74
Mean (±0.662)	6.51	10.94	13.65	14.53	11.41*
% WARE					
B		84.2	85.7	80.0	83.3
P		89.1	86.0	78.3	84.5
BR		88.4	88.0	85.9	87.4
Mean	82.4	87.2	86.6	81.4	84.4*

\* General mean