

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1964

[Full Table of Content](#)



### 64/R/AQ/C/9 Cereal Disease Reference Plots - Wheat

#### Rothamsted Research

Rothamsted Research (1965) *64/R/AQ/C/9 Cereal Disease Reference Plots - Wheat* ; Yields Of The Field Experiments 1964, pp 161 - 162 - DOI: <https://doi.org/10.23637/ERADOC-1-160>

64/C/9.1

CEREAL DISEASE REFERENCE PLOTS

(AQ)

Pennells Piece 1964, the second year

For treatments etc., see 'Results' 63/C/10. (WW = Winter wheat, SW = Spring wheat, O = Oats).

Area of each plot: 0.0180. Area harvested: Winter wheat - 0.0115, spring wheat - 0.0116.

Cultivations, etc.: Ploughed: Oct 8, 1963. All plots except winter wheat chisel ploughed: Dec 17.

Winter wheat: Seed combine drilled at 2.5 bushels: Oct 19, 1963.

'Nitro-Chalk' applied: Mar 5, 1964. Sprayed with mecoprop/2,4-D (Methoxone Extra at 7 pints in 40 gals): May 12. Combine harvested: Aug 26.

Spring wheat: Seed combine drilled at 3 bushels: Mar 11, 1964.

'Nitro-Chalk' applied: Mar 31. Sprayed with MCPA/dichlorprop (Seritox at 8 pints in 40 gals): May 15. Combine harvested: Aug 28.

Oats: Seed combine drilled at 4 bushels: Mar 7, 1964. 'Nitro-Chalk' applied: Mar 9. Sprayed with MCPA/dichlorprop (Seritox at 8 pints in 40 gals): May 15. Combine harvested: Aug 22.

Spring beans: Seed placement drilled at 200 lb (rows spaced at 10.5 ins): Feb 14, 1964. Combine harvested: Sept 2.

- NOTE: (1) Yields were taken only for winter and spring wheat.  
(2) Estimates of the incidence of take-all (*Ophiobolus graminis*), eyespot (*Cercospora herpotrichoides*) and sharp eyespot (*Corticium solani*) were made on 7 occasions for winter and 3 for spring wheat.  
(3) For previous year's results see 'Results' 63/C/10.

Standard errors per plot. Grain:

Winter wheat: 1.36 or 3.5% (5 d.f.)

Spring wheat: 2.85 or 9.4% (5 d.f.)

64/c/9.2

SUMMARY OF RESULTS

GRAIN

WINTER WHEAT

Crop in 1963

WW	0	Mean	Diff
37.9	39.1	38.2	1.2
(±0.55)	(±0.96)		(±1.11)

SPRING WHEAT

Crop in 1963

SW	0	Mean	Diff
28.0	36.6	30.2	8.6
(±1.16)	(±2.01)		(±2.32)