

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1964

[Full Table of Content](#)



### 64/W/WAS/C/16 One Year Leys for Wheat - Grass and Clover

#### Rothamsted Research

Rothamsted Research (1965) *64/W/WAS/C/16 One Year Leys for Wheat - Grass and Clover* ; Yields Of The Field Experiments 1964, pp 181 - 182 - DOI: <https://doi.org/10.23637/ERADOC-1-160>

64/C/16.1

ONE YEAR LEYS FOR WHEAT

(WAS)

A comparison of different one year leys as a preparation for wheat -  
Woburn Warren Field North - leys 1964 the first year.

Design: 3 randomised blocks of 18 plots each.

Area of each plot: 0.0212. Area harvested: 1st cut - 0.0101,  
2nd cut - 0.0138.

Treatments: All combinations of:-

1. Leys undersown in barley 1963 and cut for hay 1964, with nitrogen as follows:  
Clover: None (CO).  
Ryegrass: None (RO), 1 (R2), 2 (R4) cwt N.  
Clover-Ryegrass: None (MO) 1 (M2) cwt N.  
All N as 'Nitro-Chalk', 5/8ths to first cut, 3/8ths to second cut.
2. Potash: None (KO), 1.2 (K1), 2.4 (K2) cwt K<sub>2</sub>O as muriate of potash, 5/8ths to first cut, 3/8ths to second cut.

NOTE (1): Wheat 1965 will test in addition to the above:

3. Potash at none, 1.2, 2.4 cwt K<sub>2</sub>O
4. 'Nitro-Chalk' at none, 0.5, 1.0 cwt N.

Basal dressing. Barley nurse crop 1963: 3 cwt 21:14:14 combine drilled.  
Leys: 1.25 cwt superphosphate combine drilled.

Cultivations, etc.: Barley drilled: Apr 13, 1963. Leys undersown in barley, basal dressing applied for leys: May 6, 1963. 1st dressing of N and K applied: Mar 25, 1964. Cut: June 12. 2nd dressing of N and K applied: June 23. Cut 2nd time: Aug 5. Previous crops: Barley 1962 and 1963.

NOTE (2): Plots were sampled at each cut for estimation of dry matter and N and K per cent.

Standard errors per plot. Dry matter:

1st cut:	3.60 or 7.4% (34 d.f.)
2nd cut:	2.46 or 9.5% (34 d.f.)
Total of 2 cuts:	4.70 or 6.3% (34 d.f.)

64/C/16.2

SUMMARY OF RESULTS

GRASS, DRY MATTER

	CO	RO	R2	R4	MO	M2	Mean
1ST CUT							
(±2.08)							(±0.84)
K0	30.4	40.3	55.0	54.1	49.4	60.5	48.3
K1	35.2	42.0	61.1	59.1	50.0	54.8	50.4
K2	35.0	40.1	56.0	53.1	49.2	56.2	48.3
Mean (±1.20)	33.5	40.8	57.4	55.4	49.5	57.2	48.9
2ND CUT							
(±1.41)							(±0.58)
K0	25.8	7.1	30.8	36.2	20.0	32.9	25.5
K1	25.9	8.1	31.6	38.9	21.2	32.1	26.3
K2	27.1	6.6	31.1	40.9	18.7	33.3	26.3
Mean (±0.81)	26.3	7.3	31.2	38.6	20.0	32.8	26.0
TOTAL OF 2 CUTS							
(±2.71)							(±1.10)
K0	56.2	47.5	85.8	90.2	69.4	93.5	73.7
K1	61.1	50.1	92.7	98.0	71.2	86.9	76.7
K2	62.0	46.6	87.1	94.0	67.9	89.5	74.5
Mean (±1.56)	59.8	48.1	88.5	94.1	69.5	89.9	74.9

Mean D.M. %: 1st cut 24.3  
 2nd cut 29.6  
 Total of 2 cuts 26.9