

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1964

[Full Table of Content](#)



64/W/WAU/C/17 *Trefoil* and *Ryegrass* Green Manures - Barley

Rothamsted Research

Rothamsted Research (1965) *64/W/WAU/C/17 Trefoil and Ryegrass Green Manures - Barley* ; Yields Of The Field Experiments 1964, pp 183 - 184 - DOI: <https://doi.org/10.23637/ERADOC-1-160>

64/C/17.1

BARLEY AFTER GREEN MANURES

(WAU)

Effects of trefoil and ryegrass green manures, time of ploughing and N - Woburn Road Piece 1964 - barley following early potatoes.

Design: Single replicate of 4 x 2 x 4 in two blocks of 16 plots each, randomisation restricted.

Area of each plot: 0.0212. Area harvested: 0.0141.

Treatments. All combinations of:-

1. Green manures: None (O), trefoil (T), ryegrass (R), ryegrass sown with 0.6 cwt N as 'Nitro-Chalk' (RN).
2. Time of ploughing: Nov 26, 1963 (A), Jan 30, 1964 (S).
3. Nitrogen: None (NO), 0.3 (N1), 0.6 (N2), 0.9 (N3) cwt N as 'Nitro-Chalk'.

Basal dressing. Early potatoes: 8 cwt 17:11:22. Green manures: None. Barley: 2.5 cwt 0:20:20 combine drilled.

Cultivations, etc.: Ploughed: Sept 17 - Oct 17, 1962. Basal dressing applied for early potatoes: Apr 11, 1963. Potatoes planted: Apr 19. Earthed up: June 15. Haulm destroyed mechanically: July 23. Lifted: July 24 - Aug 2. Ground chalk applied to plots 1 - 16 at 60 cwt and to plots 17 - 32 at 20 cwt, 'Nitro-Chalk' applied to RN plots: Aug 2. Trefoil sown at 30 lb and ryegrass at 40 lb: Aug 3. A plots ploughed: Nov 26. S plots ploughed: Jan 30, 1964. 'Nitro-Chalk' treatments applied: Feb 25. Seed drilled at 2.75 bushels: Feb 28. Sprayed with mecoprop/2,4-D (Methoxone Extra at 6 pints in 40 gals): May 6. Combine harvested: Aug 21. Varieties: Trefoil - English, ryegrass - English Leafy Italian, early potatoes - Arran Pilot, barley - Maris Badger. Previous crops: Spring wheat 1961, barley 1962.

NOTE: (1) The green manures were sampled just before ploughing for estimates of dry matter and N per acre in roots plus tops.
(2) Yields were adjusted for a linear trend across the blocks and only the adjusted yields are presented.

Standard error per plot.

Barley. Grain 3.64 or 11.5% (7 d.f.)

64/C/17.2

SUMMARY OF RESULTS

GRAIN

	NO	N1	N2	N3	Mean
		(±1.29)			
Mean	22.4	29.3	35.3	39.8	31.7
		(±2.58)			(±1.29)
O	16.7	20.0	30.2	35.7	25.7
T	27.0	34.8	40.9	44.2	36.7
R	19.8	26.6	30.4	35.3	28.0
RN	26.0	36.0	39.7	43.9	36.4
		(±1.82)			(±0.91)
A	22.3	30.5	34.7	42.5	32.5
S	22.4	28.2	35.9	37.1	30.9
	O	T	R	RN	
		(±1.82)			
A	27.7	38.0	28.0	36.3	
S	23.6	35.4	28.0	36.5	

Mean D.M. %: 84.2