

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1964

[Full Table of Content](#)



64/R/RB201/DB/2 and 64/W/WB201/DB/2 Barley - Row Spacing

Rothamsted Research

Rothamsted Research (1965) *64/R/RB201/DB/2 and 64/W/WB201/DB/2 Barley - Row Spacing* ; Yields Of The Field Experiments 1964, pp 217 - 219 - DOI: <https://doi.org/10.23637/ERADOC-1-160>

64/Db/2.1

BARLEY

(RB 201 and WB 201)

Row spacing, seed rates and N - Rothamsted (R) Great Knott III and Woburn (W) Butt Close 1964.

Design (each field): 4 randomised blocks of 8 plots each, plots being split into 3 for application of N.

Area of each sub plot: 0.0056 acres.

Treatments: All combinations of:-

- Whole plots. (1) Row spacing: Seed and PK* broadcast (B): seed drilled, 4 inches between rows, PK broadcast (C): seed drilled, 7 inch rows, PK broadcast (W): seed drilled, 7 inch rows, PK combine drilled (W*).
- (2) Seed rates: 2 (L), 4 (H) bushel.
- Sub plots. (3) Nitrogen: 0.4 (N1): 0.7 (N2): 1.0 (N3) cwt N as 'Nitro-Chalk', broadcast.

* (0:20:20) to all plots - rate 2 cwt.

Cultivations, etc.:

Great Knott III (R): Ploughed twice: Sept 26 - Nov 26, 1963 and Jan 6, 1964. Seed sown: Mar 12. 'Nitro-Chalk' and broadcast PK applied: Plots 201-208 - Mar 18, remaining plots - Mar 26. Sprayed with mecoprop/2,4-D (Methoxone Extra at 6 pints in 40 gals): May 14. Combine harvested: Aug 21. Variety: Maris Badger. Previous crops: Potatoes 1962, winter wheat 1963.

Butt Close (W): Ploughed twice: Oct 2, 1963 and Jan 6, 1964. 'Nitro-Chalk' and broadcast PK applied: Feb 26. Seed sown: Mar 11. Sprayed with mecoprop/2,4-D (Methoxone Extra at 6 pints in 40 gals): May 6. Combine harvested: Aug 25. Variety: Maris Badger. Previous crops: Spring beans and carrots 1962, winter wheat 1963.

NOTES (1). Great Knott III (R): There was a drill failure on all sub plots of plot RB 205 (Treatment W*L). Estimated values were used in the analysis.

(2). Butt Close (W): Two blocks (plots WB 209 - 216 and 225 - 232) were abandoned because of poor stands and much weed growth.

Standard errors per plot. Grain:

Great Knott III (R)	Whole plot: 2.20 or 4.4% (20 d.f.)
	Sub plot: 3.00 or 6.0% (46 d.f.)
Butt Close (W)	Whole plot: 2.23 or 7.5% (7 d.f.)
	Sub plot: 4.52 or 15.1% (16 d.f.)

64/Db/2.2

SUMMARY OF RESULTS

GRAIN

ROTHAMSTED

	B	C	W	W*	Mean
	(±1.10)				(±0.55)
L	52.1	50.8	48.8	48.6	50.1
H	52.5	50.3	46.0	48.2	49.3
	(1) and (2)				(±0.53)
N1	48.4	45.7	42.8	44.3	45.3
N2	54.2	51.0	48.5	49.6	50.8
N3	54.3	54.9	51.0	51.3	52.9
Mean (±0.78)	52.3	50.5	47.4	48.4	49.7
	L	H			
	(3) and (4)				
N1	45.3	45.2			
N2	51.5	50.2			
N3	53.3	52.4			

Mean D.M. %: 85.9

(1) ±1.06 (3) ±0.75. For use in vertical and interaction comparisons.

(2) ±1.16 (4) ±0.82. For use in horizontal and diagonal comparisons.

64/Db/2.3

		GRAIN				
		WOBBURN				
		B	C	W	W*	Mean
		(±1.58)				(±0.79)
L		26.2	30.1	29.9	33.0	29.8
H		30.8	29.7	29.2	30.4	30.0
		(1) and (2)				(±1.13)
N1		20.0	22.7	19.3	21.9	21.0
N2		29.2	30.8	31.5	29.8	30.3
N3		36.3	36.2	37.9	43.5	38.5
Mean (±1.11)		28.5	29.9	29.6	31.7	29.9
		L	H			
		(3) and (4)				
N1		20.8	21.2			
N2		30.0	30.6			
N3		38.6	38.3			

Mean D.M. %: 85.4

(1) ±2.26 (3) ±1.60. For use in vertical and interaction comparisons.

(2) ±2.16 (4) ±1.53. For use in horizontal and diagonal comparisons.