

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1964

[Full Table of Content](#)



### 64/W/WP401/DD/4 Potatoes - Control of Tuber Blight

#### Rothamsted Research

Rothamsted Research (1965) *64/W/WP401/DD/4 Potatoes - Control of Tuber Blight ; Yields Of The Field Experiments 1964*, pp 233 - 234 - DOI: <https://doi.org/10.23637/ERADOC-1-160>

64/Dd/4.1

POTATOES

(WP 101)

Control of tuber blight (*Phytophthora infestans*) by fungicide sprays and haulm destruction - Woburn Horsepool 1964.

Design: 6 x 6 Latin square.

Area of each plot: 0.0360. Area harvested: 0.0141.

Treatments: No fungicide, not burnt off (0)

Fungicide sprays\*: 3 times early (E)  
4 times early (E+)  
3 times late (L).

Haulm burnt off with diquat\*\* on sprayed plots. In addition two plots per row (one control and one E+ plot on which the haulm was not burnt off) were used for sampling only.

\* 1.5 lb fungicide, containing 80% mancozeb, in 50 gals.

\*\* Reglone at 4 pints in 40 gals.

Basal dressing: 7 cwt 17:11:22.

Cultivations, etc.: Chisel ploughed: Dec 7, 1963. Ploughed: Jan 3 - 10, 1964. Basal dressing applied: Apr 14. Rotary cultivated twice: Apr 23 and 27. Potatoes machine planted: Apr 27. Earthed up: June 12. First spraying with mancozeb (E and E+ plots): June 24, second (E, E+ and L): July 14, third (E, E+ and L): July 28, final (E+ and L): Aug 10. Appropriate plots sprayed with diquat: Sept 21. Lifted: Sept 30. Variety: King Edward. Previous crops: Grass 1962, winter wheat 1963.

NOTE: Periodic samples were taken from the sample plots for weight of tubers and blight assessment in tubers.

Standard error per plot.

Total tubers: 0.681 or 6.5% (15 d.f.)

64/Da/4.2

SUMMARY OF RESULTS

0	E	E+	L	Mean
TOTAL TUBERS				
		(±0.278)		
10.30	10.21	10.52	10.66	10.42
		% WARE		
94.1	94.5	93.4	93.6	93.9

\* 1/2 lb tubers containing 500 tubers in 10 gals.  
 \*\* regions at 1 gals in 10 gals.  
 basal dressing 7 lbs 1000.  
 Observations etc.: Initial treatment Dec 7, 1957, ploughed;  
 Jan 3 - 10, 1958, basal dressing applied; Apr 10, 1958,  
 cultivated twice; Apr 27 and 31, 1958, machine ploughed;  
 Apr 27, 1958, basal dressing applied; May 12, 1958, machine  
 (E and E+ plots); June 24, 1958, (E, E+ and L); July 14, 1958,  
 (E, E+ and L); July 22, 1958, (E+ and L); Aug 10, 1958,  
 plots sprayed with Diquat; Sept 27, 1958, (E, E+ and L);  
 King Edward, previous crop: wheat 1957, winter wheat 1957.  
 BUTER: periodic samples were taken from the sample plots for weight  
 of roots and slight assessment in tubers.  
 Standard error per plot.  
 Total tuber 0.081 or 0.5% (12 d.f.).