

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1965

[Full Table of Content](#)



## 65/R/AF/C/6 Levels of N and K - Grass 8th Year

### Rothamsted Research

Rothamsted Research (1966) *65/R/AF/C/6 Levels of N and K - Grass 8th Year* ; Yields Of The Field Experiments 1965, pp 139 - 140 - DOI: <https://doi.org/10.23637/ERADOC-1-159>

65/C/6.1

GRASS

(AF)

Levels of N and K - Harwoods Piece 1965, the 8th year.

For treatments etc. see 'Results' 63/C/7 and for previous years' results see 58/Cg/2, 59/Cg/2, 60/Ci/1, 61/Dg/1, 62/C/11, 63/C/7 and 64/C/6.

Area of each plot: 0.0087. Area harvested: 0.0059.

The 1964 sowing was abandoned and the plots resown with Meadow Fescue S53 at 24 lb and Scots Timothy at 16 lb.

Cultivations, etc.: Rotary cultivated: Apr 5, 1965. First N and K dressings applied: Apr 7. Grass resown: Apr 26. Sprayed with 2,4-D butoxyethylester at 7 oz a.e. in 40 gals: June 2. Cut twice: July 21, Oct 4. N and K applied after first cut.

NOTE: Samples were taken for yield and N, P and K determinations.

Standard errors per plot. Dry matter:

1st cut:	2.38 or 9.7% (33 d.f.)
2nd cut:	4.10 or 13.3% (33 d.f.)
Total of 2 cuts:	5.34 or 9.6% (33 d.f.)

65/c/6.2

SUMMARY OF RESULTS

DRY MATTER

N	0	1	1	1	2	2	2	3	3	3	3	3	Mean
P	1	1	1	1	1	1	1	1	1	1	0	2	
K	0	0	1	2	0	1	2	0	1	2	2	2	

1ST CUT

13.7 21.4 23.5 23.4 22.1 25.6 26.2 21.9 28.3 28.2 29.2 29.7 | 24.4  
(±1.19)

Mean D.M. %: 15.7

2ND CUT

19.7 28.7 29.5 27.8 32.9 34.7 33.6 31.8 30.4 33.3 36.9 32.3 | 31.0  
(±2.05)

Mean D.M. %: 20.2

TOTAL OF 2 CUTS

33.4 50.1 53.0 51.2 55.0 60.3 59.9 53.8 58.6 61.5 66.1 62.0 | 55.4  
(±2.67)

Mean D.M. %: 18.0

Treatment symbols:

N 0 1 2 3    0.0 0.3 0.6 0.9 cwt N as 'Nitro-Chalk' 21  
 P 0 1 2    0.0 0.6 1.2 cwt P2O5 as Granular Superphosphate  
 K 0 1 2    0.0 0.3 0.6 cwt K2O as Granular Muriate of Potash.