

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1965

[Full Table of Content](#)



65/R/AP/C/8 Chemical Control of Take-all - Winter Wheat

Rothamsted Research

Rothamsted Research (1966) *65/R/AP/C/8 Chemical Control of Take-all - Winter Wheat ; Yields Of The Field Experiments 1965*, pp 143 - 144 - DOI: <https://doi.org/10.23637/ERADOC-1-159>

65/C/8.1

CHEMICAL CONTROL OF TAKE-ALL

(AP)

The chemical control of take-all (*Ophiobolus graminis*) in winter wheat - Highfield Drive 1965, the third year.

Area of each plot: 0.0072. Area harvested: 0.0071.

Treatments: None (0) and all combinations of:-

1. Rates of spraying: Heptachlor at 4 lb in 70 gals (H4), at 8 lb in 140 gals (H8).
2. Times of spraying: In 1964 only (S1), in 1963 and in 1964 (S2).

No heptachlor sprays were applied in 1965.

Basal applications: 2.5 cwt (0:20:20) combine drilled. 1 cwt N as 'Nitro-Chalk' applied as spring top dressing. Weedkiller: Mecoprop/2,4-D (Methoxone Extra at 7 pints in 40 gals).

Cultivations, etc.: Ploughed: Oct 16, 1964. Seed drilled at 160 lb: Oct 23. 'Nitro-Chalk' applied: Apr 14, 1965. Sprayed: Apr 30. Combine harvested: Aug 26. Variety: Cappelle.

- NOTES: (1) Estimates were made of the incidence of take-all and eyespot (*Cercospora herpotrichoides*) in March and June.
- (2) For previous years' results see 'Results' 63/C/9 and 64/C/8.

Standard error per plot.

Grain: 4.48 or 11.7% (8 d.f.)

65/C/8.2

SUMMARY OF RESULTS

GRAIN

0	H4 S1	H8 S1	H4 S2	H8 S2	Mean
37.4	37.8	38.1 (±2.58)	42.1	35.2	38.1

Mean D.M.%: 76.2