

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1965

[Full Table of Content](#)



65/W/WBE/C/16 *Trefoil* and Ryegrass Green Manures - Sugar Beet

Rothamsted Research

Rothamsted Research (1966) *65/W/WBE/C/16 Trefoil and Ryegrass Green Manures - Sugar Beet* ; Yields Of The Field Experiments 1965, pp 175 - 179 - DOI:

<https://doi.org/10.23637/ERADOC-1-159>

65/c/16.1

GREEN MANURES FOR SUGAR BEET

(WBE)

Effects of trefoil and ryegrass green manures and N - Woburn
Great Hill Bottom I 1965, sugar beet after barley.

Design: Two replicates of 4 x 2 x 4 in 4 blocks of 16 plots.

Area of each plot: 0.0159. Area harvested: 0.0106.

Treatments: All combinations of:-

1. Green manures undersown in barley 1964: None (O), trefoil (T), ryegrass (R), ryegrass with 0.6 cwt N as 'Nitro-Chalk' applied after barley harvest (RN).
2. Time of ploughing: Autumn (A), spring (S).
3. Nitrogen: None (NO), 0.6 (N1), 1.2 (N2), 1.8 (N3) N as 'Nitro-Chalk'.

Basal applications:

To barley: 3 cwt (17:11:22) combine drilled. Weedkiller: MCPB/MCPA (New Legumex at 5 pints in 40 gals).

To green manures: None.

To sugar beet: 5 cwt agricultural salt, 0.45 cwt K2O as muriate of potash in winter, 0.45 cwt P2O5 and 0.45 cwt K2O as (0:20:20) in seedbed.

Cultivations, etc.: Ploughed: Oct 22, 1963. Rotary cultivated twice: Mar 9 and Apr 1, 1964. Barley drilled at 140 lb: Apr 7. Trefoil undersown at 30 lb and ryegrass at 40 lb: Apr 9. Sprayed: May 15. Combine harvested: Aug 27. 'Nitro-Chalk' applied to RN plots: Sept 11. A plots ploughed: Nov 13. S plots ploughed: Feb 5, 1965. Salt and muriate of potash applied: Feb 9. Rotary cultivated: Apr 1. Rotary cultivated, basal PK compound applied: Apr 14. 'Nitro-Chalk' applied, seed drilled at 5 lb: Apr 15. Singled: May 25. Lifted: Nov 11. Varieties: Barley - Proctor, ryegrass - English Leafy Italian, trefoil - English (inoculated), sugar beet - Klein E. Previous crops: Potatoes 1961, spring wheat 1962.

NOTES: (1) Green manures were sampled for weight of dry matter and percentage N just before ploughing.

(2) There was severe frost during lifting and roots on some plots were damaged. Sugar % was estimated for two plots.

Standard errors per plot.

Roots:	1.318 or 6.3% (29 d.f.)
Total sugar:	4.94 or 6.9% (27 d.f.)
Tops:	1.200 or 7.0% (29 d.f.)

65/c/16.2

SUMMARY OF RESULTS

	NO	N1	N2	N3	Mean
ROOTS					
(±0.329)					
Mean	17.45	21.49	21.95	22.59	20.87
(±0.659)					
O	16.95	20.76	21.12	23.00	20.46
T	17.21	21.33	22.48	23.24	21.06
R	16.21	21.27	22.50	22.28	20.56
RN	19.42	22.59	21.70	21.86	21.39
(±0.466)					
A	17.74	21.77	22.85	22.41	21.19
S	17.15	21.20	21.05	22.77	20.54
(±0.466)					
(±0.233)					
(±0.466)					
A	20.76	21.34	20.78	21.88	
S	20.15	20.79	20.35	20.89	

65/C/16.3

	NO	N1	N2	N3	Mean
SUGAR %					
Mean	17.5	17.4	17.1	16.9	17.3
O	17.3	17.5	16.9	15.9	16.9
T	18.0	17.3	17.1	17.7	17.5
R	16.8	17.8	17.5	17.0	17.3
RN	18.1	17.2	17.0	16.8	17.3
A	17.5	17.3	17.0	16.9	17.2
S	17.6	17.6	17.2	16.8	17.3
	O	T	R	RN	
A	16.7	17.4	17.1	17.5	
S	17.1	17.6	17.4	17.0	

65/c/16.4

	NO	N1	N2	N3	Mean
TOTAL SUGAR					
(±1.23)					
Mean	61.0	75.0	75.1	75.9	71.7
(±2.47)					
O	58.6	72.7	71.4	73.2	69.0
T	62.2	73.7	76.8	82.0	73.7
R	53.3	75.6	78.7	74.7	70.6
RN	69.9	78.0	73.4	73.7	73.7
(±1.75)					
A	62.0	75.5	77.8	75.3	72.7
S	59.9	74.5	72.4	76.5	70.8
	O	T	R	RN	
(±1.75)					
A	69.4	74.2	70.6	76.4	
S	68.5	73.2	70.6	71.0	

65/c/16.5

	NO	N1	N2	N3	Mean
TOPS					
(±0.300)					
Mean	11.58	15.63	19.41	22.12	17.18
(±0.600)					
O	10.71	16.12	18.52	21.68	16.76
T	11.81	14.40	17.64	22.88	16.68
R	10.38	14.14	18.12	20.98	15.91
RN	13.41	17.85	23.36	22.94	19.39
(±0.424)					
A	12.30	16.26	19.35	22.58	17.62
S	10.85	15.00	19.47	21.66	16.74
	O	T	R	RN	
(±0.424)					
A	17.16	16.69	17.23	19.41	
S	16.35	16.67	14.58	19.36	