

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1965

[Full Table of Content](#)



### 65/W/WBU/C/18 Row Spacing and Fertilisers 1965-66 - Potatoes

#### Rothamsted Research

Rothamsted Research (1966) *65/W/WBU/C/18 Row Spacing and Fertilisers 1965-66 - Potatoes ; Yields Of The Field Experiments 1965*, pp 183 - 185 - DOI:

<https://doi.org/10.23637/ERADOC-1-159>

65/c/18.1

## ROW SPACING AND FERTILISERS

(WBU)

Row spacing and concentrated fertilisers - Woburn Workhouse 1965, the first year. Potatoes 1965 followed by barley 1966.

Design: Two replicates of 4 x 2 x 2 x 2 in 4 blocks of 16 plots.

Area of each plot: 0.0030. Area harvested: 0.0022.

Treatments: All combinations of:-

1. Levels of NPK (in the proportion 1.0 N, 1.0 P<sub>2</sub>O<sub>5</sub>, 1.5 K<sub>2</sub>O) to supply: None (F0), 0.66 (F1), 1.32 (F2), 2.0 (F3) cwt N.
2. Type of fertiliser: Concentrated (C), dilute (D).
3. Time of application: Before (B), after (A) rotary cultivation.
4. Spacing of setts: 14 in. x 14 in. (S14), 14 in. x 28 in. (S28).
5. Also to barley 1966, concentrated NPK fertiliser to supply none, 0.33, 0.66, 1.00 cwt N.

Basal applications: Weedkiller: diquat/paraquat (Preaglone Extra at 2 pints in 40 gals). Fungicide/insecticide: 8 lb Bordeaux mixture plus malathion/DDT (Kil at 10 fluid oz) in 40 gals, on 2 occasions. Fungicide: Bordeaux mixture at 8 lb in 40 gals on 2 occasions.

Cultivations, etc.: Ploughed: Jan 19 - 26, 1965. B fertilisers applied: Apr 13. Rotary cultivated, A fertilisers applied: Apr 21. Potatoes hand planted on the flat: Apr 21. Sprayed (weedkiller): May 7, fungicide/insecticide: July 5, July 16, fungicide: July 28, Aug 5. Sprayed with diquat (Reglone at 4 pints in 40 gals): Aug 11. Lifted: Sept 20. Variety: King Edward. Previous crops: Permanent Grass until 1964.

NOTE: The crop was severely infected with blight.

Standard error per plot.

Total tubers: 1.717 or 13.8% (30 d.f.)

65/c/18.2

SUMMARY OF RESULTS

ROW SPACING

TOTAL TUBERS

	F0	F1	F2	F3	Mean
Mean	13.39	12.46	11.69	12.16	12.43
		(±0.429)			
C		12.63	12.77	12.54	12.65
D		12.30	10.60	11.78	11.56
		(±0.607)			(±0.350)
B		12.46	11.87	11.72	12.02
A		12.46	11.51	12.60	12.19
		(±0.607)			(±0.350)
S14	14.14	13.39	13.21	12.94	13.42
S28	12.65	11.53	10.17	11.39	11.43
		(±0.607)			(±0.303)

OMITTING F0

CB	DB	CA	DA
12.56	11.48	12.73	11.65
	(±0.496)		
C S14	D S14	C S28	D S28
14.01	12.35	11.29	10.77
	(±0.496)		
E S14	A S14	E S28	A S28
12.96	13.41	11.08	10.98
	(±0.496)		

65/C/18.3

ROW SPACING

% WARE

	FO	F1	F2	F3	Mean
Mean	75.8	77.8	78.0	75.6	76.8
C		76.4	77.0	76.1	76.5
D		79.1	78.9	75.0	77.7
B		74.9	81.4	75.6	77.3
A		80.6	74.5	75.6	76.9
S14	74.2	72.2	73.9	70.9	72.8
S28	77.4	83.3	82.0	80.3	80.7

OMITTING FO

CB	DB	CA	DA
77.9	76.6	75.2	78.7
C S14	D S14	C S28	D S28
71.4	73.3	81.7	82.0
B S14	A S14	B S28	A S28
73.5	71.1	81.0	82.7