

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



# Yields of the Field Experiments 1965

[Full Table of Content](#)



## 65/R/PG/A/6 Park Grass - Hay

### Rothamsted Research

Rothamsted Research (1966) *65/R/PG/A/6 Park Grass - Hay ; Yields Of The Field Experiments 1965*, pp 25 - 26 - DOI: <https://doi.org/10.23637/ERADOC-1-159>

65/A/6.1

HAY - THE PARK GRASS PLOTS 1965

(PG)

The 110th year

For history, treatments etc., see 'Details' 1962.

Liming: All plots (excluding 5-1, 5-2, 6, 12, 15, 18-2 and all of plots 19 and 20) are divided into 4 sub-plots each, and ground chalk is applied in such a way as to establish and maintain the following reactions per sub-plot:-

Old scheme	South		North	
	Limed		Unlimed	
Sub-plot	a	b	c	d
ph existing		6		5
Plot 18-1 and 3				maximum acidity
				(a at the Eastern end).

In 1965 ground chalk was applied to sub-plots as follows (tons CaCO<sub>3</sub>):-

Plot	Sub-plot	
	b	c
1	-	2.5
4-2	1.0	4.5
9	2.0	3.5
10	1.0	4.0
11-1	5.0	4.0
11-2	3.0	4.0
13	-	1.0
18-1	-	2.0

Area of each sub-plot:\*

Plot	Area harvested
1	0.0070
2, 3	0.0088
4-1, 4-2	0.0093
11-1 and 11-2	0.0094
7, 8, 9, 10 and 13	0.0189
14, 16 and 17	0.0057
18	0.0210

Microplots: Experiments using microplots are now in progress on the following plots:-

- 5-1 and 5-2 (Effects of NPK on old grass) (see 65/22)  
6 (Effects of N levels on old grass) (see 65/22)

Cultivations, etc.: Dung applied to appropriate plots: Nov 10, 1964.

Mineral fertilisers applied: Nov 27. Ground chalk applied:  
Jan 6, 1965. Nitrogenous fertilisers applied: 1st dressing:  
Mar 10, 2nd dressing: Apr 13. Cut twice: June 28 and Dec 3.

\* The lengths harvested on sub-plots a, b, c and d varied slightly. The areas given are averages.

SUMMARY OF RESULTS

DRY MATTER

Plot No	1st cut				2nd cut				Mean	Total of 2 cuts				Total
	a	b	c	d	a	b	c	d		a	b	c	d	
1	22.8	17.0	21.9	17.9	19.9	10.0	5.1	7.3	6.9	32.8	22.1	27.1	25.2	26.8
2	20.6	22.6	23.4	21.3	21.9	6.7	5.9	7.8	6.6	27.3	28.5	31.2	27.3	28.6
3	23.4	21.9	22.1	21.9	22.3	5.7	4.9	7.0	6.2	29.1	26.8	29.2	29.0	28.5
4-1	28.0	28.5	29.0	32.6	29.5	13.5	10.7	14.3	13.5	41.5	39.2	43.3	48.2	43.1
4-2	34.9	33.9	40.0	37.5	36.6	11.6	7.2	7.4	9.7	46.5	41.1	47.5	47.2	45.6
7	44.3	45.0	41.8	34.5	41.4	17.7	17.4	21.0	18.4	62.1	62.4	62.8	51.9	59.8
8	31.1	31.4	37.8	37.1	34.3	11.1	11.0	16.3	13.7	42.2	42.3	54.1	53.6	48.1
9	51.5	47.8	50.7	56.3	51.6	19.5	13.0	6.5	8.4	11.8	71.0	60.8	57.2	64.7
10	38.5	38.2	44.0	41.9	40.6	17.8	12.8	5.8	10.7	11.8	56.2	51.0	49.8	52.6
11-1	60.9	55.4	57.4	50.5	56.1	23.8	20.3	17.6	29.5	22.8	84.8	75.8	75.1	80.0
11-2	62.3	61.7	61.6	49.9	58.9	25.3	24.6	12.5	28.1	22.6	87.6	86.3	74.1	78.0
12	28.7	27.5	28.1	19.1	19.1	13.5	13.5	16.3	16.3	47.7	47.7	41.1	41.1	44.4
13	40.2	47.4	47.3	45.8	45.2	24.0	28.6	35.1	30.3	29.5	64.2	76.0	82.4	76.1
14	43.4	50.4	46.9	42.7	45.9	10.9	11.5	16.1	17.8	14.1	54.3	61.9	63.1	60.5
15	36.7	43.4	44.4	32.7	34.7	17.0	17.0	14.5	15.7	53.7	53.7	47.2	50.4	50.4
16	43.3	43.4	44.4	36.6	41.9	15.2	16.3	20.1	16.7	17.1	58.5	59.7	64.6	53.2
17	24.2	26.0	29.8	30.3	27.6	11.4	12.1	18.7	13.3	13.9	35.6	38.1	48.4	43.6
18-1			19.5	15.2	17.4			14.5	12.8	13.7		34.1	28.0	31.0
18-2					27.2					9.7				36.9
18-3	28.6	31.6			30.1	10.8	12.1			11.4	39.4	43.7		41.5
19-1					30.1	44.4					30.1			74.5
19-2					42.1						22.6			64.7
19-3					44.7						26.7			71.3
20-1					46.6						33.9			80.5
20-2					44.9						33.9			78.8
20-3					42.1						30.3			72.4

Mean D.M. %: 1st crop: 25.7 2nd crop: 23.5

65/A/6.2