

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1965

[Full Table of Content](#)



### 65/R/RW201/DA/2 Winter Wheat and Spring Wheat - Varieties and N

#### Rothamsted Research

Rothamsted Research (1966) *65/R/RW201/DA/2 Winter Wheat and Spring Wheat - Varieties and N* ;  
Yields Of The Field Experiments 1965, pp 241 - 243 - DOI:

<https://doi.org/10.23637/ERADOC-1-159>

65/Da/3.1

# WINTER WHEAT

(RW 401 and WW 101)

Row spacing, seed rates and N - Rothamsted (R) Whittlocks and Woburn (W) Broadmead III 1965.

Design: 4 randomised blocks of 8 plots, split into 3 for N.

Area of each sub plot: 0.0045.

Treatments: All combinations of:-

Whole plots:

## 1. Row spacing etc.:

- Seed broadcast, autumn fertiliser\*\* broadcast (B)
- Seed drilled, 4 inch rows, autumn fertiliser broadcast (C)
- Seed drilled, 7 inch rows, autumn fertiliser broadcast (W)
- Seed drilled, 7 inch rows, with autumn fertiliser combine drilled. (W\*)

## 2. Seed rates:

- Whittlocks (R): 128 lb (L), 236 lb (H)
- Broadmead I (W): 139 lb (L), 231 lb (H).

Sub plots:

- 3. Nitrogen: 0.4 (N1), 0.8 (N2), 1.2 (N3) cwt N as 'Nitro-Chalk' broadcast in spring.

\*\* (6:15:15) to all plots - rate 330 lb.

Basal application: Weedkiller: Mecoprop/2,4-D (Methoxone Extra at 7 pints in 40 gals).

Cultivations, etc.:-

Whittlocks (R): Chisel ploughed: Oct 5, 1964. Seed sown, autumn fertiliser applied: Oct 27. 'Nitro-Chalk' applied: Apr 14, 1965. Sprayed: Apr 30. Combine harvested: Sept 14. Variety: Cappelle. Previous crops: Spring wheat 1963, potatoes 1964.

Broadmead III (W): Chisel ploughed: Oct 17, 1964. Autumn fertiliser applied, seed sown: Oct 23. 'Nitro-Chalk' applied: Apr 7. Sprayed: May 6. Combine harvested: Sept 15. Variety: Cappelle. Previous crops: Barley 1963, potatoes 1964.

NOTE: Whittlocks (R)

- (1) Plant counts were made on Mar 18, 1965.
- (2) Owing to a fault in the combine harvester the yields from two adjacent sub plots (WL N2 and WL N3) were bulked. Estimated yields were used in the analysis.

65/Da/3.2

Standard errors per plot. Grain:

Whittlocks (R) Whole plot: 2.24 or 5.6% (20 d.f.)  
 Sub plot: 2.13 or 5.3% (47 d.f.)  
 Broadmead III (W) Whole plot: 2.07 or 5.5% (21 d.f.)  
 Sub plot: 5.80 or 15.4% (48 d.f.)

## SUMMARY OF RESULTS

## GRAIN

## ROTHAMSTED

	B	C	W	W*	Mean
	(±1.12)				(±0.56)
L	44.3	42.7	44.0	42.8	43.5
H	37.6	36.4	35.2	36.7	36.5
	(1) and (2)				(±0.38)
N1	44.6	44.1	42.9	43.7	43.8
N2	39.4	39.2	39.9	40.2	39.7
N3	38.8	35.3	36.1	35.4	36.4
Mean (±0.79)	40.9	39.6	39.6	39.8	40.0
	L	H			
	(3) and (4)				
N1	46.4	41.2			
N2	43.6	35.8			
N3	40.4	32.5			

Mean D.M. %: 78.5

(1) ±0.75 (3) ±0.53 For use in vertical and interaction comparisons.  
 (2) ±1.00 (4) ±0.71 For use in horizontal and diagonal comparisons.

65/Da/3.3

GRAIN					
WOEBURN					
	B	C	W	W*	Mean
	(±1.04)				(±0.52)
L	42.4	38.9	37.8	39.8	39.7
H	38.0	35.5	35.3	34.1	35.7
	(1) and (2)				(±1.03)
N1	45.3	39.0	39.3	39.8	40.8
N2	40.0	37.8	39.4	35.3	38.1
N3	35.3	34.7	31.0	35.8	34.2
Mean (±0.73)	40.2	37.2	36.6	37.0	37.7
	L	H			
	(3) and (4)				
N1	41.6	40.1			
N2	41.1	35.1			
N3	36.4	32.0			

Mean D.M. %: 80.7

(1) ±2.05 (3) ±1.45 For use in vertical and interaction comparisons.  
 (2) ±1.83 (4) ±1.29 For use in horizontal and diagonal comparisons.