

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1965

[Full Table of Content](#)



### 65/R/RW401/DA/3 and 65/W/WW101/DA/3 Winter Wheat - Row Spacing

#### Rothamsted Research

Rothamsted Research (1966) *65/R/RW401/DA/3 and 65/W/WW101/DA/3 Winter Wheat - Row Spacing*  
; Yields Of The Field Experiments 1965, pp 245 - 245 - DOI:  
<https://doi.org/10.23637/ERADOC-1-159>

65/Da/1

WINTER WHEAT

(RW 301)

Varieties and nitrogen - Highfield Drive 1965.

Design: 4 randomised blocks of 6 plots each.

Area of each plot: 0.0145. Area harvested: 0.0094.

Treatments. All combinations of:-

1. Varieties: Cappelle (C), Rothwell Perdix (R).
2. Nitrogen: 0.5 (N1), 0.75 (N2), 1.00 (N3) cwt N as 'Nitro-Chalk'.

Basal applications: 280 lb (6: 15: 15) combine drilled  
Mecoprop/2,4-D (Methoxone Extra at 7 pints in 40 gals).

Cultivations, etc.: Ploughed: Oct 16, 1964. Seed drilled at  
175 lb: Oct 22. 'Nitro-Chalk' applied: Apr 13, 1965.  
Sprayed: Apr 30. Combine harvested: Aug 26. Previous  
crops: Winter wheat 1963, barley 1964.

NOTE: All plots were severely damaged by birds shortly  
before harvest.

Standard error per plot.

Grain: 3.27 or 14.1% (15 d.f.)

SUMMARY OF RESULTS

	GRAIN			Mean
	N1	N2	N3	
		(±1.63)		(±0.94)
C	22.5	24.1	23.9	23.5
R	22.5	22.6	23.4	22.8
Mean (±1.16)	22.5	23.3	23.6	23.2

Mean D.M. %: 75.2