

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1965

[Full Table of Content](#)



## 65/W/WB201/DB/2 Barley - Urea

### Rothamsted Research

Rothamsted Research (1966) *65/W/WB201/DB/2 Barley - Urea* ; Yields Of The Field Experiments 1965, pp 257 - 258 - DOI: <https://doi.org/10.23637/ERADOC-1-159>

65/Db/2.1

BARLEY

(WB 201)

Urea concentrations in NPK fertilisers - Woburn Great Hill S.W. 1965.

Design: 2 randomised blocks of 20 plots.

Area of each plot: 0.0055. Area harvested: 0.0042.

Treatments: None (0) (4 plots per block) and all combinations of:-

1. Compound fertilisers (all in the proportion N:P2O5:K2O of 2:1:1) combine drilled:-

Compounds with P as triple superphosphate:-

P, N as 100% urea.

Q, N as 66% urea and 33% ammonium nitrate.

R, N as 33% urea and 66% ammonium nitrate.

S, N as 100% ammonium nitrate.

Compounds with P and part N as monourea phosphate, remaining N as follows:-

T, as 100% urea.

U, as 66% urea and 33% ammonium nitrate.

V, as 33% urea and 66% ammonium nitrate.

W, as 100% ammonium nitrate.

K as muriate of potash in all compounds.

2. Levels: To supply 0.5 (I1), 1.0 (I2) cwt N.

Basal applications: Manures: None. Weedkiller: dichlorprop/MCPA (Cornox RK Extra at 6 pints in 50 gals).

Cultivations, etc.: Ploughed: Nov 27, 1964. Rotary cultivated: Mar 12, 1965. Seed drilled at 140 lb: Mar 15. Sprayed: May 10. Combine harvested: Aug 25. Variety: Maris Badger. Previous crops: Sugar beet 1963, barley 1964.

NOTE: Grain and straw samples were taken for yield and N percentage. Grain yields also were taken at harvest.

Standard errors per plot.

Grain: 3.91 or 10.7% (21 d.f.)

65/Db/2.2

SUMMARY OF RESULTS

GRAIN

	P	Q	R	S	T	U	V	W	Mean
	(±2.77)								(±0.98)
L1	41.9	39.1	40.7	41.4	37.7	38.3	40.6	33.9	39.2
L2	42.0	43.5	41.6	46.3	38.0	40.5	42.3	42.9	42.2
Mean (±1.96)	42.0	41.3	41.1	43.8	37.9	39.4	41.5	38.4	40.7

Plots receiving no fertiliser: 19.6 (±1.38)

General mean: 36.5

Mean D.M. %: 75.5