

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1965

[Full Table of Content](#)



### 65/R/RBE101/DC/1 Winter Beans - Irrigation and Row Spacing

#### Rothamsted Research

Rothamsted Research (1966) *65/R/RBE101/DC/1 Winter Beans - Irrigation and Row Spacing ; Yields Of The Field Experiments 1965*, pp 259 - 259 - DOI: <https://doi.org/10.23637/ERADOC-1-159>

65/Dc/1

WINTER BEANS

(RBe 101)

Row spacing, seed rates, methods of fertiliser application and irrigation - Long Hoos V 1965.

Design: A single replicate of 4 x 2 x 2 x 2 x 2 in 8 blocks of 8 plots. Irrigation on one group of 4 blocks.

Area of each plot: 0.0172.

Treatments: All combinations of:-

- Blocks: 1. Irrigation: None, 1.3 ins. water.  
Plots: 2. Row spacing: 10.5 ins. (C), 21 ins. (W).  
3. Seed rates: 275 lb (L), 412 lb (H).  
4. Fertiliser rates: 400 (F1), 560 (F2) lb compound 0:20:20, 500 (N1), 700 (N2) lb 6:15:15.  
5. Methods of fertiliser application: Broadcast (B), placed (P).

Basal applications: Manures: None. Simazine at 0.5 lb in 40 gals on 2 occasions.

Cultivations, etc.: Ploughed: Oct 15, 1964. Seed drilled, fertilisers applied: Nov 2. Sprayed: Nov 6, 1964 and Mar 15, 1965. Irrigated: June 2. Combine harvested: Sept 28. Variety: Garton's Pedigree. Previous crops: Winter and spring wheat 1963, barley 1964.

NOTE: The crop was severely attacked by Black Fly (*Aphis fabae*) and by Chocolate Spot (*Botrytis cinerea*). The experiment was therefore abandoned and no yields were taken.