Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1965



Full Table of Content

65/R//DD/8 Potatoes - Soil Fungicides

Rothamsted Research

Rothamsted Research (1966) 65/R//DD/8 Potatoes - Soil Fungicides; Yields Of The Field Experiments 1965, pp 275 - 276 - DOI: https://doi.org/10.23637/ERADOC-1-159

65/Dd/8.1

POTATOES

Soil fungicides, Long Hoos III 1965.

Design: 4 randomised blocks of 11 plots.

Area of each plot: 0.0011.

Treatments: None*(0) and all combinations of:-

1. Fungicides:

Triphenyltin acetate (A)
Tributyltin acetate (B)
Bis(triphenyltin)sulphide (C)

Triphenyltin chloride (D)

at 0.175 1b metallic Sn in 210 1b kaolin, applied to the soil and lightly forked in.

Tetrachloro-iso-phthalonitrile wettable powder (E)

sprayed into soil at 15 lb in 500 gals.

2. Times of application: June 28 - 30 (1), at first foliage blight (July 27 - 29) (2).

* 210 lb kaclin dust was applied on June 28.

Basal applications: 7 cwt (17:11:22). Weedkiller: 2 lb linuron plus 0.75 lb ion paraquat in 40 gals. Fungicide: Mancozeb at 1.2 lb in 37 gals. Insecticide: 1.5 lb phorate with seed.

Cultivations, etc.: Ploughed: Nov 23, 1964. Basal NPK applied: Apr 5, 1965. Rotary cultivated, potatoes machine planted: Apr 21. Sprayed (weedkiller): May 14, fungicide: June 30. Sprayed with undiluted BCV at 15 gals: Sept 2. Harvested: Aug 31. Variety: King Edward. Previous crops: Winter wheat 1963 and 1964.

NOTE: Tubers were examined at harvest for blight infection.

Standard error per plot.

Total tubers: 1.555 or 10.6% (30 d.f.)

65/Dd/8.2

SUMMARY OF RESULTS

TOTAL TUBERS

O Al A2 Bl B2 Cl C2 Dl D2 El E2 Mean (±0.778)

14.69 14.04 14.77 15.76 14.56 14.55 13.65 14.92 15.81 15.43 13.54 14.70