Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1965



Full Table of Content

65/W/WP701/DD/9 Potatoes - Tuber Blight

Rothamsted Research

Rothamsted Research (1966) 65/W/WP701/DD/9 Potatoes - Tuber Blight; Yields Of The Field Experiments 1965, pp 277 - 278 - DOI: https://doi.org/10.23637/ERADOC-1-159

65/Dd/9.1

POTATOES

(WP 101)

Control of tuber blight (Phytophthora infestans) by fungicide sprays and haulm destruction - Woburn Workhouse Field 1965.

Design: 6 x 6 Latin square.

Area of each plot: 0.0425. Area harvested: 0.0141.

Treatments: No fungicide (0), not burnt off.

Fungicide sprays*: 3 times early (E), burnt off with diquat**

4 times early (E+), burnt off with diquat**

3 times late (L), burnt off with diquat**

In addition two plots per row, on which haulm was not burnt off (one control and one E+ plot) were used for sampling only.

* 1.5 lb fungicide, containing 80% mancozeb, in 34 gals. ** Reglone at 4 pints in 34 gals.

Basal applications: 7 cwt (17:11:22).

Cultivations, etc.: Ploughed: Jan 19 - 26, 1965. Basal dressing applied: Apr 13. Rotary cultivated, potatoes planted: Apr 15. Earthed up: June 14. First spraying with mancozeb (E and E+ plots): July 2, second (E, E+ and L): July 15, third (E, E+ and L): July 30, final (E+ and L): Aug 13. Appropriate plots sprayed with diquat: Aug 27. Haulm destroyed mechanically: Oct 4. Lifted: Oct 13. Variety: King Edward. Previous crops: Grass 1963 and 1964.

NOTE: Periodic samples were taken from the sample plots for weight of tubers and blight assessment in tubers.

Standard error per plot.
Total tubers: 1.787 or 10.8% (15 d.f.)

				65/Dd/9.2
		SUMMARY OF RE	SULTS	
0	E	E÷	L	Mean
	T	CTAL TUBERS		
	(±0.7	730)	-1-0 170	
13.13	18.10	17.71	17.19	16.53
		% WARE		

95.1