

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1965

[Full Table of Content](#)



65/R/BK/A/1 Broadbalk - Wheat

Rothamsted Research

Rothamsted Research (1966) *65/R/BK/A/1 Broadbalk - Wheat* ; Yields Of The Field Experiments 1965, pp 9 - 11 - DOI: <https://doi.org/10.23637/ERADOC-1-159>

65/A/1.1

WHEAT - BROADBALK 1965

(BK)

For history, treatments etc. see 'Details' 1962.

Cultivations, etc.:

CROPPED SECTIONS: Ground chalk applied: Sept 9, 1964. Ploughed:
Oct 9 - 14. Dung applied: Oct 13. Autumn fertilisers
applied: Oct 19. Seed drilled at 180 lb: Oct 26. Spring
fertilisers applied: Apr 26, 1965. Second dressing of nitrate of
soda applied to plot 16: May 10. Sprayed* with MCPA at
1.13 lb and dicamba at 0.08 lb in $\frac{3}{4}$ gals: May 11. Combine
harvested: Sept 21. Variety: Squarehead's Master 13/4
(Rothamsted seed from Broadbalk field).

FALLOW SECTION: (IV) Ploughed: Oct 9 - 14, 1964, Nov 30,
July 21, 1965.

BROADBALK WILDERNESS: Cultivations, etc.:

Ungrazed meadow (north): Shrubs grubbed out: Nov 3 - 19, 1964.

Grazed meadow (centre): Grazed by sheep: Apr 26 - May 4, 1965,
May 22 - 28, June 15 - 25, July 19 - 29, Aug 18 - 31,
Oct 4 - Nov 4. Grass topped: May 3, May 28, June 25,
July 29, Sept 1.

* Except Section 5A.

65/A/1.2

SUMMARY OF RESULTS

GRAIN

Section Years after fallow	IV	VA	II	IB	VB	IA	Mean
	1	2	3	4	7	14	
2A	32.7	29.3	27.5	25.9	28.3	24.1	28.8
2B	31.5	26.3	26.6	24.2	26.9	24.9	27.5
3	23.4	13.9	8.2	8.0	7.2	9.7	13.0
5	27.1	20.6	9.3	9.0	10.7	9.7	15.7
6	24.0	14.7	14.7	13.5	13.7	12.7	16.8
7	21.5	21.7	23.1	20.9	18.9	22.2	21.6
8	26.5	24.6	23.1	26.5	22.4	19.7	24.4
9	23.3	21.4	19.2	13.5	18.6	17.7	19.6
10	18.4	22.0	18.3	14.0	14.9	10.7	17.2
11	17.9	25.8	16.5	15.5	17.1	14.7	17.8
12	18.8	21.3	17.0	17.7	14.5	16.5	17.8
13	21.2	21.1	20.0	16.4	17.4	19.3	19.6
14	21.0	20.4	18.8	19.7	17.1	14.5	19.2
15	24.7	15.3	18.3	18.5	22.1	18.7	20.2
16	20.4	22.8	20.5	20.4	21.6	16.8	20.6
17	20.3	13.5	7.9	5.6	8.7	3.7	11.5
18	24.2	15.3	18.5	15.5	23.7	19.6	20.0
19	26.1	22.9	19.1	18.4	19.6	16.9	21.3
20			12.2	17.0		12.4	14.0

Mean D.M. %: 79.3

NOTE: The yields above are calculated (as in all experiments in the 'Results') from the weight of 'first' grain delivered by the combine harvester, adjusted only for moisture content. On certain sub-plots in 1965 samples were separated into wheat, weed seeds and rubbish. Results:-

% weeds plus rubbish

Plot No

	2	5	7	9	18
Section VA	4	19	9	12	34
Section VB	2	8	3	4	2

65/A/1.3

Section Years after fallow	STRAW						Mean
	IV 1	VA 2	II 3	IB 4	VB 7	IA 14	
2A	47.3	41.1	34.3	40.9	35.9	30.9	39.6
2B	49.0	35.3	42.3	38.7	35.6	30.3	41.1
3	29.6	15.5	12.6	9.6	11.4	10.7	16.9
5	39.8	22.7	8.1	15.2	18.8	13.8	21.2
6	43.1	21.3	19.9	19.0	15.6	19.6	25.7
7	37.1	36.2	34.5	32.9	31.1	36.2	34.9
8	44.6	47.6	38.6	33.9	39.8	36.7	40.7
9	31.5	32.9	28.0	23.6	31.4	30.4	29.5
10	26.1	35.3	15.5	14.7	21.9	20.3	21.8
11	24.6	38.3	21.8	21.9	19.8	39.3	25.5
12	34.3	37.4	20.2	18.0	26.5	12.7	26.1
13	37.7	41.2	31.5	27.0	38.0	31.7	34.6
14	38.0	42.3	20.5	27.3	29.9	20.0	30.0
15	36.7	30.5	26.5	27.7	33.4	35.8	31.4
16	46.1	43.2	35.7	32.5	42.4	34.2	39.7
17	22.3	18.3	9.6	8.7	10.8	8.2	14.0
18	40.5	22.8	24.5	22.1	39.2	28.0	30.4
19	36.3	31.9	25.7	25.5	28.5	24.4	29.6
20			14.3	21.3		27.9	19.2

Mean D.M.%: 88.3