

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1966

[Full Table of Content](#)



66/W/WBT/C/4 Methods of Application of Fertiliser 1965-66 - Spring Wheat

Rothamsted Research

Rothamsted Research (1967) *66/W/WBT/C/4 Methods of Application of Fertiliser 1965-66 - Spring Wheat* ; Yields Of The Field Experiments 1966, pp 131 - 133 - DOI:

<https://doi.org/10.23637/ERADOC-1-158>

66/c/4.3

Plots receiving no fertiliser in 1965

GRAIN				STRAW			
FO	F1	F2	Mean	FO	F1	F2	Mean
20.3	35.8 (±1.61)	45.6	33.9	9.4	23.6	33.8	22.3

General mean 37.4

25.6

Mean D.M. %: 84.8

79.8

Errata to 'Results' 65/c/4.3

Plots receiving no fertiliser in 1964. Means should read:
 Whittlocks (R): 30.7 not 31.2
 Broadmead (W): 32.0 not 30.7

66/C/5.1

GRASS

(AF)

Levels of N and K - Harwoods Piece 1966, the 9th year.

For treatments etc. see 'Results' 63/C/7.1 and 65/C/6.2 and for previous years' results see 58/Cg/2, 59/Cg/2, 60/C1/1, 61/Dg/1, 62/C/11, 63/C/7, 64/C/6 and 65/C/6.

Area of each plot: 0.0087. Area harvested: 1st cut - 0.0057, 2nd, 3rd, 4th cuts - 0.0059.

Cultivations, etc.: N, P and K fertilisers applied: Mar 28, 1966. Cut 4 times: May 17, June 28, Aug 15, Oct 13. N and K applied after first 3 cuts.

NOTE: Crop samples were taken for N, P and K determinations.

Standard errors per plot. Dry matter:

1st cut:	2.07 or 7.5% (33 d.f.)
2nd cut:	1.70 or 6.8% (33 d.f.)
3rd cut:	1.90 or 10.3% (33 d.f.)
4th cut:	1.37 or 11.5% (33 d.f.)
Total of 4 cuts:	4.78 or 5.8% (33 d.f.)