

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1966

[Full Table of Content](#)



66/R/AZ/C/8 Row Spacing N and Paraquat - Lucerne 3rd Year

Rothamsted Research

Rothamsted Research (1967) *66/R/AZ/C/8 Row Spacing N and Paraquat - Lucerne 3rd Year* ; Yields Of The Field Experiments 1966, pp 143 - 147 - DOI: <https://doi.org/10.23637/ERADOC-1-158>

66/c/8.3

DRY MATTER

2ND CUT

	NO	N1	N2	Mean
Mean (± 0.52)	26.4	26.1 (± 0.74)	26.2	26.3 (± 0.43)
C	27.9	26.5	26.6	27.0
W	24.9	25.8 (± 0.74)	25.9	25.5 (± 0.43)
PO	26.5	25.4	25.5	25.8
PI	26.2	26.9	26.9	26.7
	C	W		
	(± 0.60)			
PO	26.4	25.2		
PI	27.5	25.8		

Mean D.M. %:16.3

66/c/8.4

DRY MATTER

3RD CUT

	NO	N1	N2	Mean
Mean (± 0.39)	15.0	16.4 (± 0.55)	15.3	15.6 (± 0.32)
C	15.8	16.2	15.2	15.7
W	14.2	16.5 (± 0.55)	15.4	15.4 (± 0.32)
PO	15.8	17.1	16.4	16.4
PI	14.2	15.7	14.3	14.7
	C	W		
	(± 0.45)			
PO	16.7	16.2		
PI	14.8	14.6		

Mean D.M. %: 23.8

66/c/8.5

DRY MATTER
TOTAL OF 3 CUTS

	NO	N1	N2	Mean
Mean (± 1.13)	73.0	75.3 (± 1.60)	76.9	75.0 (± 0.92)
C	76.3	76.4	77.2	76.6
W	69.7	74.2 (± 1.60)	76.6	73.5 (± 0.92)
PO	77.1	80.8	82.1	80.0
PI	68.8	69.8	71.7	70.1
	C	W		
		(± 1.31)		
PO	81.4	78.6		
PI	71.8	68.4		

Mean D.M. %: 20.2

Table with multiple columns and rows, containing numerical data and text. The table is very faint and difficult to read, but appears to be a data table with several columns and rows. The text is mirrored and bleed-through from the reverse side of the page.

66/C/9.1

IRRIGATION

(IRA and IRB)

The effect of irrigation on potatoes and barley - Great Field I and II 1966. For previous years' results see 'Results' 65/C/14.

Design:

Potatoes: Two replicates of 4 x 2 x 3 x 2 in two blocks. Each block is divided into 4 whole plots for irrigation treatments, and at right angles into 6 strips for varieties and spacing.

There is a further split for N.

Barley: Two randomised blocks of 4 plots, with plots split for N.

Area of each sub plot:

Potatoes: 0.0214
Barley: 0.1286

Area harvested:

0.0080
0.0689

Treatments: Potatoes: All combinations of:-

Whole plots. 1. Irrigation: None (O), early (A), late (B), full (C).

To strips of 1/6th plots.

2. Varieties: Majestic (M), King Edward (E).
Unchitted seed.

3. Spacing: Seed 12 (S1), 15 (S2), 18 inches (S3) apart in rows.

1/12th plots. 4. Nitrogen: 1.2 (N1), 1.8 (N2) cwt N as compound fertiliser (basal) and 'Nitro-Chalk'.

Barley: All combinations of:-

Main plots. 1. Irrigation: None (O), early (A), late (B), full (C).

Sub plots. 2. Nitrogen: 0.6 (N1), 0.8 (N2) cwt N as compound fertiliser (basal) and 'Nitro-Chalk'.

Basal applications:

Potatoes: 12 cwt compound fertiliser (10:10:18). Weedkiller: Linuron at 1 lb and paraquat at 0.75 lb ion in 37 gals.

Fungicide: Mancozeb at 1.2 lb in 37 gals on 3 occasions.

Barley: 3 cwt compound fertiliser (20:10:10) combine drilled.

Weedkiller: Mecoprop/2,4-D (Methoxone Extra at 6 pints in 40 gals).

Cultivations, etc.:-

Potatoes: Sprayed with dalapon at 8.8 lb a.e. in 40 gals:

Nov 8, 1965. Ploughed: Jan 5, 1966. Basal compound