Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1966



Full Table of Content

66/R/AZ/C/8 Row Spacing N and Paraquat - Lucerne 3rd Year

Rothamsted Research

Rothamsted Research (1967) 66/R/AZ/C/8 Row Spacing N and Paraquat - Lucerne 3rd Year; Yields Of The Field Experiments 1966, pp 143 - 147 - DOI: https://doi.org/10.23637/ERADOC-1-158

66/c/8.3

DRY MATTER

2ND CUT

	NO	Nl	N2	Mean
Mean (±0.52)	26.4	26.1	26.2	26.3
		(±0.74)		(±0.43)
C W	27.9 24.9	26.5 25.8	26.6 25.9	27.0 25.5
		(±0.74)		(±0.43)
PO Pl	26.5 26.2	25.4 26.9	25. 5 26. 9	25.8 26.7
	C	W		
	(±0,60)			
P0 P1	26.4 27.5	25.2 25. 8		

Mean D.M. %:16.3

66/c/8.4

DRY MATTER

3RD CUT

	NO	Nl	N2	Mean
Mean (±0.39)	15.0	16.4	15.3	15.6
	3,85	(±0.55)		(±0.32)
C W	15.8 14.2	16.2 16.5	15.2 15.4	15.7 15.4
	8.88	(±0.55)		(±0.32)
PO Pl	15.8 14.2	17.1 15.7	16.4 14.3	16.4 14.7
	c	W		
	(±0.45)			
P0 P1	16.7 14.8	16.2 14.6		

Mean D.M. %: 23.8


66/c/8.5

DRY MATTER

TOTAL OF 3 CUTS

	NO	Nl	N2	Mean
Mean (±1.13)	73.0	75.3	76.9	75.0
		(±1.60)		(±0.92)
C ₩	76.3 69.7	76.4 74.2	77.2 76.6	76.6 73.5
		(±1.60)		(±0.92)
PO Pl	77.1 68.8	80.8 69.8	82.1 71.7	80.0 70.1
	С	W		
	(±1.31)			
PO Pl	81.4 71.8	78.6 68.4		

Mean D.M. %: 20.2



66/c/9.1

IRRIGATION

(IRA and IRB)

The effect of irrigation on potatoes and barley - Great Field I and II 1966. For previous years' results see 'Results' 65/C/14.

Design:

Potatoes: Two replicates of 4 x 2 x 3 x 2 in two blocks. Each block is divided into 4 whole plots for irrigation treatments, and at right angles into 6 strips for varieties and spacing. There is a further split for N.

Barley: Two randomised blocks of 4 plots, with plots split for N.

Area of each sub plot: Potatoes: 0.0214 Barley: 0.1286 Area harvested: 0.0080 0.0689

Treatments: Potatoes: All combinations of:-

Whole plots. 1. Irrigation: None (0), early (A), late (B), full (C).

To strips of 1/6th plots.

 Varieties: Majestic (M), King Edward (E). Unchitted seed.

Spacing: Seed 12 (S1), 15 (S2), 18 inches (S3).
apart in rows.

1/12th plots. 4. Nitrogen: 1.2 (N1), 1.8 (N2) cwt N as compound fertiliser (basal) and 'Nitro-Chalk'.

Barley: All combinatons of:-

Main plots. 1. Irrigation: None (0), early (A), late (B), full (C).

Sub plots. 2. Nitrogen: 0.6 (N1), 0.8 (N2) cwt N as compound fertiliser (basal) and 'Nitro-Chalk'.

Basal applications:

Potatoes: 12 cwt compound fertiliser (10:10:18). Weedkiller: Linuron at 1 lb and paraquat at 0.75 lb ion in 37 gals. Fungicide: Mancozeb at 1.2 lb in 37 gals on 3 occasions. Barley: 3 cwt compound fertiliser (20:10:10) combine drilled. Weedkiller: Mecoprop/2,4-D (Methoxone Extra at 6 pints in

40 gals).

Cultivations, etc.:-

Potatoes: Sprayed with dalapon at 8.8 lb a.e. in 40 gals: Nov 8, 1965. Ploughed: Jan 5, 1966. Basal compound