

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1966

[Full Table of Content](#)



66/R/PGM/C/13 N P K - Old Grass (Park Grass Plot 5 Microplots)

Rothamsted Research

Rothamsted Research (1967) *66/R/PGM/C/13 N P K - Old Grass (Park Grass Plot 5 Microplots)* ;
Yields Of The Field Experiments 1966, pp 165 - 173 - DOI:

<https://doi.org/10.23637/ERADOC-1-158>

66/c/13.3

PLOT 5/1: DRY MATTER

2ND CUT

Excluding K1 and K6 plots

	P0	P1	P2	P4	Mean
Mean (± 1.42)	16.7	21.3	26.1	28.3	23.1
		(± 2.84)			(± 1.42)
K0	10.3	18.1	20.8	23.0	18.0
K2	18.1	21.2	30.5	29.1	24.7
K4	20.6	25.2	26.1	29.0	25.2
K8	18.0	20.7	27.0	32.1	24.5
		(± 2.01)			(± 1.00)
N1	17.9	20.7	24.5	26.7	22.4
N2	15.6	21.8	27.8	29.9	23.8
	K0	K2	K4	K8	
		(± 2.01)			
N1	18.6	23.1	23.8	24.2	
N2	17.4	26.3	26.6	24.8	
	K1* and K6* plots				
	K1	K6	Mean		
	(± 2.84)		(± 2.01)		
N1	18.0	19.1	18.5		
N2	18.5	18.0	18.2		
Mean (± 2.01)	18.2	18.5	18.4		

* Applied 1965

General mean: 22.2

Mean D.M. %: 21.5

66/C/13.4

PLOT 5/1: DRY MATTER

3RD CUT

Excluding K1 and K6 plots

	P0	P1	P2	P4	Mean
Mean (± 0.86)	18.0	22.5	24.0	26.2	22.7
		(± 1.72)			(± 0.86)
K0	16.6	21.8	22.8	20.2	20.4
K2	17.9	23.8	22.4	26.7	22.7
K4	18.4	21.6	24.8	29.0	23.4
K8	19.2	22.8	26.2	28.9	24.3
		(± 1.22)			(± 0.61)
N1	17.3	21.8	22.7	26.6	22.1
N2	18.7	23.3	25.4	25.9	23.3
	K0	K2	K4	K3	
		(± 1.22)			
N1	19.4	22.9	23.3	22.7	
N2	21.3	22.5	23.6	25.9	
	K1* and K6* plots				
	K1	K6	Mean		
	(± 1.72)		(± 1.22)		
N1	16.4	19.8	18.1		
N2	18.7	22.4	20.6		
Mean (± 1.22)	17.6	21.1	19.4		

* Applied 1965

General mean: 22.0

Mean D.M. %: 19.5

66/C/13.5

PLOT 5/1: DRY MATTER

TOTAL OF 3 CUTS

Excluding K1 and K6 plots

	P0	P1	P2	P4	Mean
Mean (± 2.23)	41.1	64.2	72.6	77.4	63.8
		(± 4.46)			(± 2.23)
K0	33.1	59.5	63.2	63.2	54.8
K2	43.9	65.5	77.1	78.9	66.4
K4	44.8	69.4	75.1	82.7	68.0
K8	42.4	62.4	74.9	84.8	66.2
		(± 3.16)			(± 1.58)
N1	42.7	61.7	66.3	73.4	61.0
N2	39.4	66.8	78.9	81.4	66.6
	K0	K2	K4	K8	
		(± 3.16)			
N1	53.8	63.5	63.3	63.4	
N2	55.7	69.2	72.7	68.9	
	K1* and K6* plots				
	K1	K6	Mean		
		(± 4.46)	(± 3.16)		
N1	49.4	56.7	53.0		
N2	53.7	55.1	54.4		
Mean (± 3.16)	51.5	55.9	53.7		

* Applied 1965

General mean: 61.8

Mean D.M. %: 21.1

66/c/13.6

PLOT 5/2: DRY MATTER

1ST CUT

Excluding K1 and K6 plots

	P0	P1	P2	P4	Mean
Mean (± 1.10)	35.5	33.9	39.0	33.6	35.5
		(± 2.19)			(± 1.10)
K0	34.4	32.9	34.5	34.7	34.1
K2	36.4	32.5	37.5	34.5	35.2
K4	35.2	36.5	43.2	32.0	36.7
K8	36.0	33.5	41.0	33.0	35.9
		(± 1.55)			(± 0.78)
N1	29.1	24.8	28.4	24.4	26.7
N2	41.8	42.9	49.7	42.7	44.3
	K0	K2	K4	K8	
		(± 1.55)			
N1	26.1	28.4	26.8	25.4	
N2	42.1	42.0	46.7	46.3	
	K1* and K6* plots				
	K1	K6	Mean		
		(± 2.19)	(± 1.55)		
N1	26.5	29.4	28.0		
N2	37.2	38.5	37.8		
Mean (± 1.55)	31.8	33.9	32.9		

* Applied 1965

General mean: 35.0

Mean D.M. %: 19.1

66/c/13.7

PLOT 5/2: DRY MATTER

2ND CUT

Excluding K1 and K6 plots

	P0	P1	P2	P4	Mean
Mean (± 1.01)	28.3	27.1	26.1	29.3	27.7
		(± 2.03)			(± 1.01)
K0	27.7	28.4	31.6	26.5	28.5
K2	29.2	26.4	26.7	32.8	28.8
K4	27.6	25.4	24.2	27.5	26.2
K8	28.6	28.4	22.0	30.4	27.4
		(± 1.43)			(± 0.72)
N1	28.3	25.9	27.8	27.5	27.4
N2	28.2	28.4	24.5	31.1	28.0
	K0	K2	K4	K8	
		(± 1.43)			
N1	27.4	29.1	26.4	26.8	
N2	29.7	28.5	26.0	27.9	
	K1* and K6* plots				
	K1	K6	Mean		
		(± 2.03)	(± 1.43)		
N1	26.5	29.3	27.9		
N2	33.7	33.1	33.4		
Mean (± 1.43)	30.1	31.2	30.7		

* Applied 1965

General mean: 28.3

Mean D.M. %: 20.7

66/c/13.8

PLOT 5/2: DRY MATTER

3RD CUT

Excluding K1 and K6 plots

	P0	P1	P2	P4	Mean
Mean (± 0.78)	27.5	26.6	27.1	29.0	27.6
		(± 1.57)			(± 0.78)
K0	26.0	27.5	26.8	27.6	27.0
K2	27.8	26.8	29.1	30.2	28.5
K4	28.4	25.1	28.1	28.1	27.4
K8	27.7	27.2	24.5	30.1	27.4
		(± 1.11)			(± 0.56)
N1	27.1	23.7	25.7	25.3	25.5
N2	27.8	29.5	28.6	32.7	29.7
	K0	K2	K4	K8	
		(± 1.11)			
N1	25.6	27.1	2.8	24.3	
N2	28.3	29.8	30.0	30.5	
	K1* and K6* plots				
	K1	K6	Mean		
		(± 1.57)	(± 1.11)		
N1	26.8	27.2	27.0		
N2	30.5	32.1	31.3		
Mean (± 1.11)	28.6	29.7	29.1		

* Applied 1965

General mean: 27.9

Mean D.M. %: 16.3

66/C/13.9

PLOT 5/2: DRY MATTER

TOTAL OF 3 CUTS

Excluding K1 and K6 plots

	P0	P1	P2	P4	Mean
Mean (± 2.17)	91.2	87.6	92.3	91.9	90.8
		(± 4.33)			(± 2.17)
K0	88.1	88.7	92.8	88.8	89.6
K2	93.3	85.7	93.3	97.6	92.5
K4	91.2	87.0	95.4	87.7	90.3
K8	92.3	89.1	87.6	93.6	90.6
		(± 3.06)			(± 1.53)
N1	84.5	74.5	81.9	77.2	79.5
N2	97.9	100.8	102.7	106.6	102.0
	K0	K2	K4	K8	
		(± 3.06)			
N1	79.0	84.6	78.0	76.5	
N2	100.2	100.3	102.7	104.7	
	K1* and K6* plots				
	K1	K6	Mean		
		(± 4.33)	(± 3.06)		
N1	79.7	85.9	82.8		
N2	101.3	103.7	102.5		
Mean (± 3.06)	90.5	94.8	92.7		

* Applied 1965

General mean: 91.1

Mean D.M. %: 18.7

TABLE 1
SUMMARY OF DATA

(continued)

Year	1970	1971	1972	1973	1974	1975
Production (1000 tons)	1.00	1.10	1.20	1.30	1.40	1.50
Consumption (1000 tons)	1.00	1.10	1.20	1.30	1.40	1.50
Stock (1000 tons)	0.00	0.00	0.00	0.00	0.00	0.00
Imports (1000 tons)	0.00	0.00	0.00	0.00	0.00	0.00
Exports (1000 tons)	0.00	0.00	0.00	0.00	0.00	0.00
Balance (1000 tons)	0.00	0.00	0.00	0.00	0.00	0.00
Price (1000 tons)	1.00	1.10	1.20	1.30	1.40	1.50
Cost (1000 tons)	1.00	1.10	1.20	1.30	1.40	1.50
Profit (1000 tons)	0.00	0.00	0.00	0.00	0.00	0.00
Loss (1000 tons)	0.00	0.00	0.00	0.00	0.00	0.00
Net (1000 tons)	0.00	0.00	0.00	0.00	0.00	0.00
Revenue (1000 tons)	1.00	1.10	1.20	1.30	1.40	1.50
Cost (1000 tons)	1.00	1.10	1.20	1.30	1.40	1.50
Profit (1000 tons)	0.00	0.00	0.00	0.00	0.00	0.00
Loss (1000 tons)	0.00	0.00	0.00	0.00	0.00	0.00
Net (1000 tons)	0.00	0.00	0.00	0.00	0.00	0.00

* Applied 1970
 General 1970
 from 1970 to 1975

66/c/14.1

PARK GRASS MICROPLOTS

(PGM 41-80)

Plot 6, 1966, the second year

For details of treatments etc., and for previous year's results see 'Results' 65/C/33.

Cultivations, etc.: P, K, Na and Mg fertilisers applied:

Dec 16, 1965. 'Nitro-Chalk' applied: Mar 7, 1966. Mecoprop treatment applied at 45 oz a.e. in 50 gals: First spraying: Apr 26, second spraying at same rate: July 28. Cut: May 4, June 1, July 8, Aug 1, Sept 8, Oct 7. 'Nitro-Chalk' applied after every cut except the last.

Standard error per plot.

Dry matter, total of all cuts: 4.09 or 5.5% (27 d.f.)