

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1966

[Full Table of Content](#)



66/R/HWF/A/3 Hoosfield - Wheat After Fallow

Rothamsted Research

Rothamsted Research (1967) *66/R/HWF/A/3 Hoosfield - Wheat After Fallow* ; Yields Of The Field Experiments 1966, pp 17 - 18 - DOI: <https://doi.org/10.23637/ERADOC-1-158>

66/A/3.1

WHEAT AFTER FALLOW - HOOSFIELD 1966

(HWF)

For history, treatments, etc. see 'Details' 1962.

The winter wheat was damaged by wheat bulb fly (*Leptohylemia coarctata*) and plots A3 and A4 were resown with spring wheat. Plot A1, which was less severely attacked, was left in winter wheat.

Area of each plot: Cappelle (C) - 0.0552, Rothwell Perdix (R) - 0.0690, Kloka (K) - 0.124. Area harvested: 0.0368.

Cultivations, etc.:-

Cropped plots: Ploughed: Oct 14, 1965. Seed drilled at 190 lb: Nov 5. Sections A3 and A4 rotary cultivated and redrilled with Kloka spring wheat at 230 lb: May 2, 1966. Winter wheat sprayed with ioxynil/mecoprop (Actril C at 6 pints in 40 gals): May 16. Spring wheat sprayed with MCPA at 1.5 lb a.e. in 40 gals: June 16. Winter wheat combine harvested: Sept 3. Spring wheat combine harvested: Sept 7. Fallowed plots: Ploughed 3 times: Oct 14, 1965, May 23, 1966 and July 15.

66/A/3.2

SUMMARY OF RESULTS

Plot	A3		A4		A1	
	1		1		3	
No of years of fallow						
Strip	A	B	A	B	A	B
Variety	K	K	K	K	C	R
GRAIN						
	10.0	9.0	10.2	9.6	11.2	13.7
STRAW						
	8.0	6.5	8.1	7.1	4.6	7.0

Mean D.M. %: Grain: 79.7
 Straw: 84.8