

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1966

[Full Table of Content](#)



66/R/BP/C/17 Legumes and Barley - Cereals and Hay

Rothamsted Research

Rothamsted Research (1967) *66/R/BP/C/17 Legumes and Barley - Cereals and Hay* ; Yields Of The Field Experiments 1966, pp 181 - 183 - DOI: <https://doi.org/10.23637/ERADOC-1-158>

66/c/17.3

OATS

GRAIN

Undersown 1965	Undersown 1966				Mean
	-	T	N1	N2	
			(+1.87)*		
-	43.2	36.8	36.5	43.5	40.0
T	46.9	37.6	38.1	46.4	42.3
Undersown 1966			(+1.87)*		
-			42.1	48.0	45.1
T			32.5	41.9	37.2
Mean (± 1.32)			37.3	45.0	41.1

* For use in horizontal and interaction comparisons only.

Mean D.M. %: 80.0

	SPRING BEANS Grain	HAY: DRY MATTER			Total of 3 cuts
		1st cut	2nd cut	3rd cut	
Mean	31.1	23.8	29.0	28.6	81.5
Mean D.M. %:	72.0	17.5	16.5	16.9	17.0

DATE

CHAIN

Year	1987	1988	1989	1990
Mean (±1.5)	21.1	21.1	21.1	21.1
1987	21.1	21.1	21.1	21.1
1988	21.1	21.1	21.1	21.1
1989	21.1	21.1	21.1	21.1
1990	21.1	21.1	21.1	21.1

* For use in horizontal and vertical comparisons only.

Mean Date: 21.1

DATE

CHAIN

Year	1987	1988	1989	1990
Mean (±1.5)	21.1	21.1	21.1	21.1
1987	21.1	21.1	21.1	21.1
1988	21.1	21.1	21.1	21.1
1989	21.1	21.1	21.1	21.1
1990	21.1	21.1	21.1	21.1

66/c/18.1

PREVIOUS CROPS & N FOR BARLEY

(BQ)

The effect of previous cropping and nitrogen on the yield of barley - Stackyard 1966, the second year - barley.

Design: 3 randomised blocks of 9 plots, split into 3 for N.

Area of each sub plot: 0.0096. Area harvested: 0.0064.

Treatments: All combinations of:-

Whole plots (applied 1965):-

1. Cropping: Spring wheat (W), Kale (K), Italian Ryegrass (G).
2. Nitrogen: None (R0), 1.0 (R2), 2.0 cwt (R4) N as 'Nitro-Chalk'.

Sub plots (applied to barley 1966):-

3. Nitrogen: None (N0), 0.5 (N1), 1.0 cwt (N2) as 'Nitro-Chalk'.

Basal applications: 2.5 cwt (0:20:20) combine drilled. Weed-killer: Mecoprop/2,4-D (Methoxone Extra at 6 pints in 30 gals).

Cultivations, etc.: Ploughed: Jan 28, 1966. Seed drilled at 145 lb: Mar 10. 'Nitro-Chalk' applied: Mar 11. Sprayed: May 17. Combine harvested: Aug 20. Variety: Maris Badger.

NOTE: For the previous year's results see 'Results' 65/C/27.

Standard errors per plot. Grain:

Whole plot: 3.04 or 11.0% (16 d.f.)

Sub plot: 3.07 or 11.1% (36 d.f.)