

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1966

[Full Table of Content](#)



66/R/BY/C/19 Previous Crops and N for Barley 1965-66 - Spring Wheat, Kale and Ryegrass

Rothamsted Research

Rothamsted Research (1967) *66/R/BY/C/19 Previous Crops and N for Barley 1965-66 - Spring Wheat, Kale and Ryegrass* ; Yields Of The Field Experiments 1966, pp 189 - 191 - DOI:
<https://doi.org/10.23637/ERADOC-1-158>

66/c/19.3

RYEGRASS

DRY MATTER

NO	N2	N4	Mean
1ST CUT			
10.6	(±1.73) 31.2	38.9	26.9
2ND CUT			
16.7	(±1.11) 23.8	34.8	25.1
3RD CUT			
8.7	(±0.34) 9.4	10.2	9.4
TOTAL OF 3 CUTS			
36.0	(±2.51) 64.3	84.0	61.4

Mean D.M. %: 1st cut: 18.1
 2nd cut: 19.6
 3rd cut: 16.6
 Total of 3 cuts: 18.1

2010-2011

STATE OF CALIFORNIA

DEPARTMENT OF REVENUE

SALES TAX

Year	Rate	Amount	Percentage	Total
2010	8.25	1,200,000,000	100%	1,200,000,000
2011	8.25	1,200,000,000	100%	1,200,000,000
2012	8.25	1,200,000,000	100%	1,200,000,000
2013	8.25	1,200,000,000	100%	1,200,000,000
2014	8.25	1,200,000,000	100%	1,200,000,000
2015	8.25	1,200,000,000	100%	1,200,000,000

1,200,000,000

1,200,000,000

1,200,000,000

1,200,000,000

1,200,000,000

1,200,000,000

66/c/20.1

BARLEY FOLLOWED BY RYEGRASS

(EQ)

The rate of action of P fertilisers, Sawyers II 1966, the second year.

Design: 4 replicates of 7 x 2 x 2 plus 2 plots per block without P, 8 blocks of 16 plots.

Area of each plot:

Barley: 0.0018
Ryegrass: 0.0018

Area harvested:

0.0009
0.0007

Treatments, to barley only: No P (0) (2 plots per block) and all combinations of:-

1. Phosphatic fertilisers:

Triple superphosphate	(S)
Potassium metaphosphate	(K)
Triple superphosphate plus potassium metaphosphate*	(SK)
Magnesium ammonium phosphate	(M)
Triple superphosphate plus magnesium ammonium phosphate*	(SM)
Potassium metaphosphate plus magnesium ammonium phosphate*	(KM)
Triple superphosphate plus potassium metaphosphate plus magnesium ammonium phosphate*	(SKM)

* Each material supplying half the P (one third in treatment SKM).

2. Levels of P: 12 lb (I1), 24 lb (I2) P.

3. Type of fertiliser: Powder, less than 1 mm. (P), granular 1-4 mm. (G).

All applied to barley with 'Nitro-Chalk', muriate of potash and kieserite in amounts adjusted to give a total of 60 lb N, 100 lb K and 25 lb Mg.

Basal applications: To ryegrass: 60 lb N and 50 lb K (16:0:16) on 2 occasions. Weedkillers: To barley: Ioxynil and mecoprop (Actril C at 4 pints in 50 gals), paraquat to stubble at 1 lb ion in 50 gals.

Cultivations, etc.

Barley: Ploughed: Jan 8, 1966. Fertilisers applied, seed drilled at 160 lb: Mar 10. Weedkiller applied: May 19. Harvested green: June 24. Paraquat applied: June 29. Variety: Maris Badger.