

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1966

[Full Table of Content](#)



66/W/C/25 K and Protein Synthesis - Grass

Rothamsted Research

Rothamsted Research (1967) *66/W/C/25 K and Protein Synthesis - Grass* ; Yields Of The Field Experiments 1966, pp 213 - 216 - DOI: <https://doi.org/10.23637/ERADOC-1-158>

66/C/25.1

GRASS

Effect of K on protein synthesis - Woburn Stackyard Series C 1966,
2nd year.

Design: 4 randomised blocks of 10 plots, with N on blocks for
3rd and 4th cuts.

Area of each plot: 0.0014. Area harvested: 0.0008.

Treatments: All combinations of:-

Blocks (to 3rd and 4th cuts only):

1. N in addition to basal: None, 100 lb N as ammonium nitrate on June 21, repeated Aug 2 (i.e. for 3rd and 4th cuts).

Plots:

2. Species: Cocksfoot S37 (C), Meadow Fescue S215 (M), sown 1965.
3. Levels of K: None (K0), 60 (K1), 120 (K2), 180 (K3), 240 lb K (K4) as sulphate of potash applied after the second cut.

Basal application: 100 lb N as ammonium nitrate in spring and
after the first, second and third cuts.

Cultivations etc.: N applied: Mar 9, May 10, June 21, Aug 2, 1966.

K applied: June 23. Cut four times: May 10, June 21, Aug 2,
Sept 27.

NOTES: (1) After each cut samples were analysed for total N, protein
N and K. After the last cut, samples were taken to
assess the residual effects.

(2) For the previous year's results see 'Results' 65/C/32.

Standard errors per plot. Dry matter:

1st cut:	1.21 or 5.2% (27 d.f.)
2nd cut:	1.77 or 8.8% (27 d.f.)
Total of 1st and 2nd cuts:	1.74 or 4.0% (27 d.f.)
3rd cut:	1.53 or 7.6% (18 d.f.)
4th cut:	1.77 or 6.6% (18 d.f.)
Total of 3rd and 4th cuts:	3.01 or 6.4% (18 d.f.)

66/C/25.2

SUMMARY OF RESULTS

DRY MATTER

	K0	K1	K2	K3	K4	Mean
1ST CUT						
			(± 0.61)			(± 0.27)
C	14.3	16.2	18.0	18.5	20.1	17.4
M	28.1	28.8	30.0	29.7	30.8	29.5
Mean (± 0.43)	21.2	22.5	24.0	24.1	25.4	23.5
2ND CUT						
			(± 0.88)			(± 0.40)
C	19.1	21.4	20.4	20.2	21.5	20.5
M	20.1	20.4	19.5	18.1	19.1	19.5
Mean (± 0.62)	19.6	20.9	19.9	19.1	20.3	20.0
TOTAL OF 1ST AND 2ND CUTS						
			(± 0.87)			(± 0.39)
C	33.4	37.6	38.4	38.6	41.6	37.9
M	48.2	49.3	49.6	47.8	49.9	48.9
Mean (± 0.61)	40.8	43.5	44.0	43.2	45.7	43.4
Mean D.M. %: 1st cut: 19.2 2nd cut: 22.8 1st and 2nd cuts: 21.0						

66/C/25.3

DRY MATTER

	K0	K1	K2	K3	K4	Mean
3RD CUT						
			(±0.77)*			
N1	16.4	18.8	20.9	20.0	21.5	19.5
N2	16.9	20.7	21.1	21.9	23.3	20.8
			(±0.77)			(±0.34)
C	15.3	19.2	21.0	21.8	22.9	20.1
M	18.0	20.4	21.0	20.2	21.9	20.3
Mean (±0.54)	16.7	19.8	21.0	21.0	22.4	20.2

4TH CUT						
			(±0.89)*			
N1	22.5	25.4	27.9	27.9	28.9	26.5
N2	20.9	26.4	27.3	29.9	29.8	26.9
			(±0.89)			(±0.40)
C	19.9	24.9	27.2	28.8	29.8	26.1
M	23.5	26.8	28.0	29.1	28.9	27.3
Mean (±0.63)	21.7	25.9	27.6	28.9	29.4	26.7

Mean D.M. %: 3rd cut: 23.3
4th cut: 25.0

* For use in horizontal and interaction comparisons only

66/C/25.4

DRY MATTER

	K0	K1	K2	K3	K4	Mean
TOTAL OF 3RD AND 4TH CUTS						
			(±1.51)*			
N1	38.9	44.2	48.8	47.9	50.5	46.1
N2	37.8	47.2	48.4	51.8	53.1	47.7
			(±1.51)			(±0.67)
C	35.3	44.2	48.2	50.5	52.8	46.2
M	41.4	47.2	49.1	49.3	50.8	47.6
Mean (±1.07)	38.3	45.7	48.6	49.9	51.8	46.9

Mean D.M. %: 24.1

* For use in horizontal and interaction comparisons only