


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1966

[Full Table of Content](#)



---

### 66/R/CA/C/28 Levels and Forms of N - Spring Beans

#### Rothamsted Research

Rothamsted Research (1967) *66/R/CA/C/28 Levels and Forms of N - Spring Beans* ; Yields Of The Field Experiments 1966, pp 223 - 224 - DOI: <https://doi.org/10.23637/ERADOC-1-158>

66/C/28.1

SPRING BEANS

(CA/Be)

Rates and forms of N for beans followed by wheat - Great Knott II 1966, the first year.

Design: 3 randomised blocks of 12 plots.

Area of each plot: 0.0193. Area harvested: 0.0121.

Treatments: None (3 plots per block) and all combinations of:-

1. Forms of nitrogen: Ammonium nitrate (A), nitrate of soda (N), sulphate of ammonia (S).
2. Rates of N: None (NO), 1 (N1), 2 (N2), 3 (N3) cwt N, broadcast.

Basal applications: 360 lb (0:14:28) placement drilled.

Insecticide: Demeton-s-methyl (Metasystox at 12 fluid oz in 37 gals).

Cultivations, etc.: Ploughed: Nov 16, 1965. Seed drilled at 200 lb: Mar 8, 1966. Nitrogen applied: Mar 14. Insecticide applied: June 14. Combine harvested: Sept 23. Variety: Pedigree Tick. Previous crops: Winter wheat and barley 1964, barley 1965.

NOTE: Plant samples were taken for nodulation counts and grain samples for 1000 grain weight and % N.

Standard error per plot.

Grain: 3.00 or 7.3% (21 d.f.)

66/c/28.2

SUMMARY OF RESULTS

GRAIN

	NO	N1	N2	N3	Mean
		(±1.73)			(±1.00)
A		40.2	42.4	42.6	41.7
N		41.3	42.6	42.3	42.1
S		39.3	39.5	43.9	40.9
Mean (±1.00)	39.8	40.3	41.5	42.9	41.1*

\* General mean

Mean D.M. %: 75.8