

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1966

[Full Table of Content](#)



66/R/RW101/DA/1 and 66/W/WW101/DA/1 Winter Wheat - Row Spacing

Rothamsted Research

Rothamsted Research (1967) *66/R/RW101/DA/1 and 66/W/WW101/DA/1 Winter Wheat - Row Spacing* ; Yields Of The Field Experiments 1966, pp 243 - 246 - DOI:
<https://doi.org/10.23637/ERADOC-1-158>

66/Da/1.1

WINTER WHEAT

(RW101 and WW101)

Row spacing, seed rates and N - Rothamsted (R) Great Knott I and Woburn (W) Workhouse 1966.

Design: 2 replicates of a 4 x 2 x 4 factorial arranged in 4 blocks of 8 whole plots, with N on half plots. (4 - 3 + 2 - 1 on whole plots Great Knott I (R), 3 - 2 + 1 - 0 on Workhouse (W)).

Area of each sub plot: 0.0069. Area harvested: 0.0069.

Treatments: All combinations of:-

Whole plots.

1. Row spacing etc.:

Seed broadcast, autumn fertiliser** broadcast (B)

Seed drilled, 4 inch rows, autumn fertiliser broadcast (C)

Seed drilled, 7 inch rows, autumn fertiliser broadcast (W)

Seed drilled, 7 inch rows, with autumn fertiliser combine drilled (W*)

2. Seed rates: 140 lb (L), 240 lb (H).

Sub plots:

3. Nitrogen in spring:

Great Knott I (R): 0.4 (N1), 0.8 (N2), 1.2 (N3), 1.6 (N4) cwt N as 'Nitro-Chalk' broadcast.

Workhouse (W): None (NO), 0.4 (N1), 0.8 (N2), 1.2 (N3) cwt N as 'Nitro-Chalk' broadcast.

** Great Knott I (R): (6:15:15) to all plots - rate 340 lb.

Workhouse (W): (8:20:16) to all plots - rate 310 lb.

NOTE: At Rothamsted because of a mistake in an instruction nitrogen was applied to the wrong plots: in order to achieve balance certain additions were made making the total applications listed.

Basal applications: Great Knott I (R): Ioxynil/MCPA (Actril A at 2 pints in 40 gals).

66/Da/1.2

Cultivations, etc.:

Great Knott I (R): Deep-tine cultivated twice: Nov 3 and 4, 1965.
Seed sown, NPK applied: Nov 5. 'Nitro-Chalk' applied: Apr 13, 1966,
completed: Apr 21. Sprayed: Apr 29. Combine harvested: Aug 24.
Variety: Cappelle. Previous crops: Barley 1964, potatoes 1965.
Workhouse (W): Deep-tine cultivated: Oct 27, 1965. Seed sown,
NPK applied: Nov 3. 'Nitro-Chalk' applied: Apr 27, 1966.
Combine harvested: Aug 27. Variety: Cappelle. Previous crops:
Grass 1964, potatoes 1965.

Standard errors per plot. Grain:

Great Knott I (R).	Whole plot: 2.57 or 5.3% (14 d.f.)
	Sub plot: 3.11 or 6.4% (16 d.f.)
Workhouse (W).	Whole plot: 2.88 or 5.2% (14 d.f.)
	Sub plot: 3.98 or 7.1% (16 d.f.)

66/Da/1.3

SUMMARY OF RESULTS

GREAT KNIGHT I (R)

GRAIN

	B	C	W	W*	Mean
Mean (± 0.91)	49.2	48.7	48.9	46.4	48.3
	(± 1.29)				(± 0.64)
L	49.3	50.6	50.9	48.3	49.8
H	49.0	46.8	47.0	44.5	46.8
	(1) and (2)				(3) and (4)
N1	49.9	49.8	47.3	47.0	48.5
N2	52.0	51.3	50.7	48.1	50.5
N3	46.4	48.1	49.4	45.0	47.2
N4	48.4	45.6	48.3	45.4	47.0
	L	H			
	(5) and (6)				
N1	49.6	47.4			
N2	50.9	50.2			
N3	48.7	45.7			
N4	49.9	44.0			

- (1) (± 1.56) (5) (± 1.10) For use in vertical and interaction comparisons involving N4-N2 or N3-N1
 (2) (± 1.69) (6) (± 1.20) For use in all other comparisons
 (3) (± 0.78) For use in comparisons N4-N2 or N3-N1
 (4) (± 0.85) For use in all other comparisons

66/Da./1.4

WORKHOUSE (W)

GRAIN

	B	C	W	W*	Mean
Mean (± 1.02)	55.9	56.7	55.4	55.3	55.8
		(± 1.44)			(± 0.72)
L	56.5	57.7	56.1	54.9	56.3
H	55.4	55.8	54.6	55.6	55.3
		(1) and (2)			(3) and (4)
NO	52.8	55.3	53.7	53.0	53.7
N1	57.4	60.1	59.8	57.5	58.7
N2	59.6	58.8	56.2	56.7	57.8
N3	53.9	52.6	52.0	53.8	53.1
	L	H			
		(5) and (6)			
NO	52.4	55.0			
N1	57.8	59.6			
N2	59.3	56.4			
N3	55.8	50.3			

- (1) (± 1.99) (5) (± 1.41) For use in vertical and interaction comparisons involving N3-N1 or N2-NO
- (2) (± 2.02) (6) (± 1.43) For use in all other comparisons
- (3) (± 1.00) For use in comparisons N3-N1 or N2-NO
- (4) (± 1.01) For use in all other comparisons