

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1966

[Full Table of Content](#)



66/R/RW601/DA/4 and 66/W/WW301/DA/4 Spring Wheat - Anhydrous Ammonia

Rothamsted Research

Rothamsted Research (1967) *66/R/RW601/DA/4 and 66/W/WW301/DA/4 Spring Wheat - Anhydrous Ammonia* ; Yields Of The Field Experiments 1966, pp 251 - 253 - DOI:

<https://doi.org/10.23637/ERADOC-1-158>

66/Da/4.1

SPRING WHEAT

(RW 601 and WW 301)

Anhydrous ammonia as a fertiliser - Rothamsted (R) Great Knott III and Woburn (W) Lansome 1966.

Design: 4 randomised blocks of 8 plots.

Area of each plot: 0.0321. Area harvested: Great Knott III (R) - 0.0208, Lansome (W) - 0.0207.

Treatments: All combinations of:-

1. Forms of N: Broadcast 'Nitro-Chalk' (B), injected anhydrous ammonia (I).
2. Levels of N: 0.5 (N1), 1.0 (N2), 1.5 (N3) cwt N together with no nitrogen - without (NO), with (NOI), the injector running empty through the soil.

Basal applications: 280 lb (0:20:20). Weedkiller: Great Knott III (R): Mecoprop/2,4-D (Methoxone Extra at 6 pints in 40 gals). Lansome (W): Ioxynil/mecoprop (Actril C at 5 pints in 35 gals).

Cultivations, etc.:

Great Knott III (R): Ground chalk applied at 22 cwt: Nov 8, 1965. Ploughed: Nov 15, 1965. Anhydrous ammonia injected: Mar 2, 1966. Seed drilled at 170 lb, basal PK and 'Nitro-Chalk' applied: Mar 16, 1966. Sprayed: May 17. Combine harvested: Sept 7. Variety: Kloka. Previous crops: Barley 1964, spring beans 1965.

Lansome (W): Ploughed: Sept 30 - Oct 25, 1965. Anhydrous ammonia injected: Mar 3, 1966. Seed drilled at 160 lb, basal PK and 'Nitro-Chalk' applied: Mar 15. Sprayed: May 13. Combine harvested: Sept 6. Variety: Kloka. Previous crops: Winter wheat 1964, barley 1965.

Standard errors per plot. Grain:

Great Knott III (R): 2.15 or 5.3% (21 d.f.)
Lansome (W): 3.57 or 14.8% (21 d.f.)

66/Da/4.2

SUMMARY OF RESULTS
GREAT KNOTT III (R)

GRAIN				
	N1	N2	N3	Mean
		(±1.07)		(±0.62)
B	43.7	45.6	43.6	44.3
I	36.6	46.3	46.9	43.3
Mean (±0.76)	40.2	46.0	45.3	43.8

	NO	NOI	Mean
	31.7	31.1	31.4
	(±1.07)		

General mean: 40.7

Mean D.M. %: 83.0

66/Da/4.3

LANSOME (W)

GRAIN

	N1	N2	N3	Mean
		(±1.78)		(±1.03)
B	23.9	32.8	37.2	31.3
I	17.7	30.2	36.5	28.1
Mean (±1.26)	20.8	31.5	36.9	29.7

	NO	NOI	Mean
	6.6	7.6	7.1
	(±1.78)		

General mean: 24.1

Mean D.M. %: 85.2