

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1966

[Full Table of Content](#)



66/R/RB101/DB/1AND 66/W/WB101/DB/1 Barley - Row Spacing

Rothamsted Research

Rothamsted Research (1967) *66/R/RB101/DB/1AND 66/W/WB101/DB/1 Barley - Row Spacing* ; Yields Of The Field Experiments 1966, pp 263 - 265 - DOI: <https://doi.org/10.23637/ERADOC-1-158>

66/Db/1.1

BARLEY

(RB101 and WB101)

Row spacing, seed rates and N - Rothamsted (R) Whittlocks N.E. and Woburn (W) Horsepool E. 1966.

Design: 2 replicates of 4 x 2 x 4 factorial arranged in 4 blocks of 8 whole plots, with N on half plots. (3 - 2 + 1 - 0 on whole plots.)

Area of each sub-plot: 0.0069. Area harvested: 0.0069.

Treatments: All combinations of:-

Whole plots:

1. Row spacing etc.:

- Seed broadcast, PK** broadcast (B)
- Seed drilled, 4 inch rows, PK broadcast (C)
- Seed drilled, 7 inch rows, PK broadcast (W)
- Seed drilled, 7 inch rows, with PK combine drilled (W*)

2. Seed rates: 110 lb (L), 220 lb (H).

Sub-plots:

3. Nitrogen: None, 0.4, 0.7, 1.0 cwt N as 'Nitro-Chalk'.

** (0:20:20) to all plots - rate 220 lb.

Basal applications: Weedkiller: Mecoprop/2,4-D (Methoxone Extra at 6 pints in 36 gals).

Cultivations, etc.:

Whittlocks N.E. (R): Ploughed: Nov 1-9, 1965. Seed sown, fertilisers applied: Mar 10, 1966. Sprayed: May 11. Combine harvested: Aug 19. Variety: Maris Badger. Previous crops: Potatoes 1964, winter wheat 1965.

Horsepool E. (W): Ploughed: Oct 28, 1965. PK fertiliser applied and seed sown: Mar 11, 1966. 'Nitro-Chalk' applied: Mar 14. Sprayed: May 13. Combine harvested: Aug 19. Variety: Maris Badger. Previous crops: Potatoes 1964, barley 1965.

Standard errors per plot. Grain:

Whittlocks N.E. (R)	Whole plot: 2.10 or 4.6% (14 d.f.)
	Sub plot: 2.18 or 4.7% (16 d.f.)
Horsepool E. (W)	Whole plot: 2.10 or 4.4% (14 d.f.)
	Sub plot: 3.67 or 7.7% (16 d.f.)

66/D6/1.2

SUMMARY OF RESULTS

WHITFLOCKS N.E. (R)

GRAIN

	B	C	W	W*	Mean
Mean (± 0.74)	45.8	45.9	47.1	45.7	46.1
	(± 1.05)				(± 0.53)
L	46.1	45.8	48.8	46.3	46.8
H	45.6	45.9	45.4	45.0	45.5
	(1) and (2)				(3) and (4)
NO	36.2	37.6	39.8	36.9	37.6
N1	48.1	47.8	49.5	48.2	48.4
N2	50.6	48.9	49.9	48.3	49.4
N3	48.5	49.2	49.3	49.3	49.0
	L	H			
	(5) and (6)				
NO	39.4	35.8			
N1	47.9	48.9			
N2	50.5	48.3			
N3	49.2	48.8			

- (1) (± 1.84) and (5) (± 1.30) For use in vertical and interaction comparisons involving N3-N1 or N2-NO
- (2) (± 1.30) and (6) (± 0.92) For use in all other comparisons
- (3) (± 0.55) For use in comparisons N3-N1 or N2-NO
- (4) (± 0.65) For use in all other comparisons

66/Db/1.3

HORSEPOOL E. (W)

GRAIN

	B	C	W	W*	Mean
Mean (± 0.74)	47.2	47.6	48.1	46.7	47.4
	(± 1.05)				(± 0.52)
L	47.2	49.1	50.1	47.1	48.4
H	47.3	46.1	46.1	46.4	46.5
	(1) and (2)				(3) and (4)
N0	46.0	43.3	45.6	42.6	44.4
N1	48.6	51.3	49.8	48.4	49.5
N2	51.4	49.0	49.5	49.7	49.9
N3	43.0	47.0	47.7	46.3	46.0
	L	H			
	(5) and (6)				
N0	44.4	44.4			
N1	50.8	48.2			
N2	52.2	47.6			
N3	46.3	45.7			

- (1) (± 1.84) and (5) (± 1.30) For use in vertical and interaction comparisons involving N3-N1 or N2-N0
 (2) (± 1.67) and (6) (± 1.18) For use in all other comparisons
 (3) (± 0.92) For use in comparisons N3-N1 or N2-N0
 (4) (± 0.83) For use in all other comparisons