

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1966

[Full Table of Content](#)



66/W/WP101/DD/9 Potatoes - Varieties

Rothamsted Research

Rothamsted Research (1967) *66/W/WP101/DD/9 Potatoes - Varieties* ; Yields Of The Field Experiments 1966, pp 293 - 294 - DOI: <https://doi.org/10.23637/ERADOC-1-158>

66/Da/9.1

POTATOES

(WP 101)

Varieties - Woburn Butt Close 1966.

Design: 4 x 4 Latin square.

Area of each plot: 0.0096. Area harvested: 0.0048.

Treatments:-

Varieties: Pentland Dell (D), King Edward (E), Majestic (see note below), Maris Piper (P).

Basal applications, etc.: 10 tons dung, 7 cwt (17:11:22).

Fungicide: Mancozeb 1.2 lb in 33 gals on 3 occasions. Weedkiller: Linuron at 1 lb a.i., and paraquat 0.75 lb ion in 37 gals.

Cultivations, etc.: Deep-tine cultivated: Sept 16, 1965. Ploughed: Nov 15. Dung applied: Jan 11, 1966. Ploughed: Jan 28. Basal NPK applied: Mar 23. Rotary cultivated and planted: Mar 31. Earthed up (rotoridged): May 4. Sprayed weedkiller: May 7. Sprayed fungicide: June 29, July 18, Aug 4. Haulm destroyed mechanically: Sept 3. Lifted: Sept 13. Previous crops: Barley 1964, fallow 1965.

NOTE: The seed of the varieties D, E and P was once-grown ex Rothamsted, H certificate. Majestic was also included but once-grown seed was not available so a Scotch SS stock was used. Because of fungus diseases this gave only 60% of normal plants (the other varieties gave full plant populations) and was therefore excluded from the analysis. The mean yield of total tubers for Majestic was 9.55 tons, mean % ware 87.7.

Standard errors per plot.

Total tubers: 1.792 or 17.1% (6 d.f.)

66/Dd/9.2

SUMMARY OF RESULTS

D	E	P	Mean
TOTAL TUBERS			
	(±0.896)		
7.72	10.71	13.04	10.49
% WARE			
72.7	69.8	84.2	75.5

NOTE: The seed of the varieties D, E and F was raised from the tubers of the variety D. The seed of the variety E was raised from the tubers of the variety E. The seed of the variety F was raised from the tubers of the variety F. The tubers of the variety D were raised from the seed of the variety D. The tubers of the variety E were raised from the seed of the variety E. The tubers of the variety F were raised from the seed of the variety F. The tubers of the variety D were raised from the seed of the variety D. The tubers of the variety E were raised from the seed of the variety E. The tubers of the variety F were raised from the seed of the variety F.