


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/W/WCW/B/8 Cultivation-weedkiller Rotation

Rothamsted Research

Rothamsted Research (1968) *67/W/WCW/B/8 Cultivation-weedkiller Rotation ; Yields Of The Field Experiments 1967*, pp 105 - 108 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/B/8.1

CULTIVATION - WEEDKILLER ROTATION

(WCW)

A comparison of weed control by various cultivation methods and by a pre-emergence weedkiller - Woburn Great Hill I and II 1967, the eighth year.

For history, rotation, treatments etc., to barley, see 'Results' 60/B/11, 61/B/11, 62/B/11, 63/B/11, 64/B/10, 65/B/9 and 66/B/8.

Area of each plot: 0.0482. Area harvested: Potatoes - 0.0107, barley - 0.0227.

Potatoes.

Treatments: All combinations of:-

1. Primary cultivations: Ploughed (P), rotary cultivated (R), deep-tine cultivated (T).
2. Weedkiller: None, normal cultivations (M), linuron at 0.25 lb, plus paraquat at 0.75 lb ion in 50 gals (S).

NOTE: Weeds grew very vigorously and so all plots were rotary-ridged, except plot 36 where the haulm was exceptionally large and would have been damaged.

Basal applications:

Barley: 340 lb (20:10:10) combine drilled. Weedkiller: Ioxynil/mecoprop (Actril C at 5 pints in 25 gals).

Potatoes: 10 cwt (17:11:22). Weedkiller: Dalapon at 9 lb in 33 gals. Fungicide: Mancozeb at 1.2 lb in 30 gals. Fentin acetate at 4.2 oz plus maneb at 1.4 oz in 30 gals. Haulm destroyer: Undiluted BOV at 15 gals.

Cultivations, etc.:

Potatoes: Weedkiller applied: Oct 7, 1966. Ground chalk applied at 40 cwt: Nov 3. T plots deep-tine cultivated (two strokes): Nov 15. P plots ploughed: Nov 15-16. R plots rotary cultivated: Nov 21. P and T plots spring-tine cultivated, and basal NPK applied: Mar 20, 1967. P and T plots spring-tine cultivated (twice): Mar 22. R plots rotary cultivated, potatoes machine planted: Mar 23. M plots harrowed: Apr 20. M plots re-ridged: Apr 28. S plots sprayed with weedkiller: May 3. M plots earthed up: May 10. All plots (except plot 36) rotary ridged:

67/B/8.2

June 9. Sprayed fungicide: July 1 and July 26. Sprayed haulm destroyer: Aug 22. Lifted: Sept 21. Variety: Maris Piper. Barley: Ground chalk applied at 40 cwt: Nov 3, 1966. T plots deep-tine cultivated (two strokes), and P plots ploughed: Nov 15. R plots rotary cultivated: Nov 21. All plots spring-tine cultivated: Feb 15, 1967. 'Nitro-Chalk' applied: Feb 16. Seed combine drilled at 140 lb: Mar 3. Sprayed weedkiller: Apr 28. Combine harvested: Aug 18. Variety: Maris Badger.

Standard errors per plot.

Potatoes, total tubers: 1.371 or 32.9% (8 d.f.)
Barley, grain: 1.73 or 6.0% (8 d.f.)

67/B/8.3

SUMMARY OF RESULTS

POTATOES

	M	S	Mean
TOTAL TUBERS			
	(±0.970)	(±0.685)	(±0.560)
P	4.67	4.30	4.42
R	3.84	4.51	4.29
T	3.33	4.01	3.79
Mean (±0.560)	3.95	4.28	4.17

% WARE

P	80.5	82.5	81.8
R	79.3	81.4	80.7
T	77.6	83.5	81.5
Mean	79.1	82.5	81.4

67/B/8.4

BARLEY

GRAIN

	N1	N2	N3	Mean
		(±1.23)		(±0.71)
P	26.1	29.0	30.1	28.4
R	26.5	29.3	34.1	29.9
T	24.1	29.1	32.0	28.4
Mean (±0.71)	25.6	29.1	32.0	28.9

Mean D.M. %: 83.7