


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/R/AQ/C/5 Cereal Disease Reference Plots - Wheat

Rothamsted Research

Rothamsted Research (1968) *67/R/AQ/C/5 Cereal Disease Reference Plots - Wheat* ; Yields Of The Field Experiments 1967, pp 133 - 134 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/C/5.1

CEREAL DISEASE REFERENCE PLOTS

(AQ)

Pennells Piece 1967, the fifth year

For treatments etc., see 'Results' 63/C/10 (WW = Winter wheat, SW = Spring wheat, O = Oats, Be = Spring beans).

The variety of oats is now Manod, and the variety of beans is now called Tarvin.

Area of each plot: 0.0180. Area harvested: Winter wheat - 0.0115, spring wheat - 0.0116.

Cultivations, etc.: Sprayed with aminotriazole at 4 lb and ammonium thiocyanate at 3.7 lb in 40 gals: Sept 17, 1966. Ploughed: Oct 20.

Winter wheat: Seed combine drilled at 160 lb: Oct 27, 1966.

'Nitro-Chalk' applied: Mar 20. Sprayed with ioxynil/mecoprop (Actril C at 6 pints in 32 gals): Apr 27, 1967. Combine harvested: Aug 30.

Spring wheat: Seed combine drilled at 175 lb: Mar 4, 1967.

'Nitro-Chalk' applied: Mar 20. Sprayed with ioxynil/mecoprop (Actril C at 5 pints in 20 gals): May 12. Combine harvested: Aug 23.

Oats: Seed combine drilled at 160 lb: Mar 8, 1967. 'Nitro-

Chalk' applied: Mar 20. Sprayed with ioxynil/mecoprop (Actril C at 5 pints in 20 gals): May 12. Combine harvested: Aug 17.

Spring beans: Seed placement drilled at 200 lb: Mar 8, 1967.

Sprayed twice with demeton-s-methyl (Metasystox at 12 fluid oz in 37 gals, and by hand at 15 fluid oz in 37 gals): June 13, July 7. Combine harvested: Sept 1.

- NOTES: (1) Yields were taken for winter and spring wheat only. (Crop Sequences 2, 3, 4, and 6).
(2) Estimates were made throughout the growing season of the incidence of take-all (*Ophiobolus graminis*) and eyespot (*Cercospora herpotrichoides*).
(3) For previous years' results see 'Results' 63/C/10, 64/C/9, 65/C/9 and 66/C/7.

67/c/5.2

SUMMARY OF RESULTS

GRAIN

Crop	1963	1964	1965	1966	Mean
	W	W	Be	W	
	W	Be	O	W	
	Be	O	W	W	
	O	W	W	W	

WINTER WHEAT

49.7	45.7	36.6	37.0	42.3
------	------	------	------	------

SPRING WHEAT

47.9	39.8	36.5	38.4	40.6
------	------	------	------	------

Mean D.M. %: Winter wheat: 84.4
Spring wheat: 81.8