


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/R/BH/C/11 Sod Seeding and Pests - Winter Wheat

Rothamsted Research

Rothamsted Research (1968) *67/R/BH/C/11 Sod Seeding and Pests - Winter Wheat* ; Yields Of The Field Experiments 1967, pp 159 - 160 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/C/11.1

WINTER WHEAT

(BH)

Sod seeding and pests, New Zealand 1967, the third year.

Design: 3 randomised blocks of 3 plots, split into 2.

Area of each sub plot: 0.0226. Area harvested: 0.0152.

Treatments: All combinations of:-

Whole plots: 1. Seedbed preparation: Ploughed 1965, 1966 and 1967 (M). Ploughed with ioxynil spray 1965, ploughed 1966 and 1967 (MI). Seeded direct each year, with paraquat 1965 and 1966, aminotriazole* and paraquat** 1967 (P).

Sub plots: 2. Insecticide spray: None (O), sprayed with DDT,*** diazinon*** and chlordane*** (D).

* At 4 lb in 40 gals (with ammonium thiocyanate at 3.7 lb).

** At 2 lb ion in 40 gals.

*** A mixture of DDT at 9.6 lb, diazinon at 4.8 lb and chlordane at 9.6 lb, in 160 gals.

Basal applications: 24 cwt ground chalk, 340 lb (6:15:15), 450 lb 'Nitro-Chalk' in spring. Weedkiller: Mecoprop/2,4-D (Methoxone Extra at 6 pints in 20 gals).

Cultivations, etc.: M and MI plots ploughed: Sept 15, 1966. Aminotriazole and ammonium thiocyanate applied: Sept 16. Ground chalk applied: Sept 23. Insecticides and paraquat applied: Oct 21. Basal NPK compound applied: Oct 24. Spring-tine cultivated M and MI treatments: Oct 25. Seed drilled at 180 lb: Oct 26. Basal 'Nitro-Chalk' applied: Apr 18, 1967. Weedkiller applied: May 9. Combine harvested: Aug 30. Variety: Cappelle.

NOTES: (1) Counts of soil fauna were made in June, August and September. Earthworm counts were included in September.

(2) For previous years' results see 'Results' 65/C/24 and 66/C/15.

Standard errors per plot. Grain:

Whole plot: 2.32 or 6.4% (4 d.f.)

Sub plot: 2.15 or 6.0% (6 d.f.)

67/C/11.2

SUMMARY OF RESULTS

	GRAIN			
	M	MI	P	Mean
	(1) and (2)			(±0.72)
O	42.4	42.4	24.9	36.6
D	39.0	42.6	25.1	35.6
Mean (±1.34)	40.7	42.5	25.0	36.1

Mean D.M. %: 83.8

- (1) (±1.60) For use in horizontal and diagonal comparisons
 (2) (±1.24) For use in vertical and interaction comparisons