

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1967

[Full Table of Content](#)



---

### 67/R/BP/C/13 Legumes and Barley - Barley

#### Rothamsted Research

Rothamsted Research (1968) *67/R/BP/C/13 Legumes and Barley - Barley* ; Yields Of The Field Experiments 1967, pp 163 - 165 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/C/13.1

LEGUMES AND BARLEY

(BP)

Effects of crop sequences and green manures - Stackyard 1967, the third year - barley.

Design: 2 randomised blocks of 10 plots, plots being split into 4 for N.

Area of each sub plot: 0.0101. Area harvested: 0.0062.

Treatments: All combinations of:-

Whole plots:

1. Crop sequences (C):

	1	2	3	4	5	6	7	8	9	10
1965	B	B	B	B	B	B	B	B	B	B
Undersown	Cl	T	T	-	-	-	-	-	T	T
1966	H	B	O	B	O	B	O	Be	B	O
Undersown	-	T	T	T	T	-	-	-	-	-
1967	B	B	B	B	B	B	B	B	B	B

Half plots:

2. Nitrogen (applied 1965 and 1966): 0.4 (R1), 0.8 (R2) cwt N as 'Nitro-Chalk' in seedbed (none to beans and hay).

Quarter plots:

3. Nitrogen to barley 1967: None (N0), 0.3 (N1), 0.6 (N2), 0.9 (N3) cwt N as 'Nitro-Chalk' in seedbed.

B = barley, H = hay, O = oats, Be = spring beans, Cl = red clover, T = trefoil.

Basal applications: 280 lb (0:20:20) combine drilled: Weedkiller: Mecoprop/2,4-D (Methoxone Extra at 6 pints in 35 gals).

Cultivations, etc.: Hay plots ploughed: Sept 15, 1966. Corrective K applied to hay plots at 1.0 cwt K<sub>2</sub>O as muriate of potash: Oct 4. Ploughed: Jan 13, 1967. Seed drilled at 140 lb: Mar 17. 'Nitro-Chalk' applied: Mar 23. Sprayed: May 11. Combine harvested: Aug 21. Variety: Maris Badger.

NOTE: For previous years' results see 'Results' 65/C/26 and 66/C/17.

67/C/13.2

Standard errors per plot. Grain

Crop Sequences 2 3 4 5 6 7 9 10  
Whole plot: 9.75 or 38.2% (6 d.f.)  
1/4 plot: 4.57 or 17.9% (14 d.f.)

Crop Sequences 1 and 8  
1/4 plot: 3.11 or 9.7% (6 d.f.)

67/c/13.3

SUMMARY OF RESULTS

GRAIN

Crop Sequences 2 3 4 5 6 7 9 10

	R1	R2	NO	N1	N2	N3	Mean
			(1) and (2)				(±4.88)
C2	18.5	19.9	16.1	19.0	20.0	21.8	19.2
C3	34.8	26.1	18.4	27.9	31.4	44.1	30.4
C4	24.2	23.4	10.1	25.1	28.3	31.7	23.8
C5	33.2	27.2	18.7	28.8	38.8	34.5	30.2
C6	23.2	19.6	14.2	16.1	20.8	34.4	21.4
C7	29.7	30.6	17.1	33.5	32.7	37.2	30.2
C9	21.3	15.2	9.2	17.4	22.5	24.1	18.3
C10	30.5	31.4	19.5	23.9	41.0	39.4	30.9
Mean	26.9	24.2	15.4	24.0	29.4	33.4	25.5
			(±1.14)				

Mean D.M. %: 81.1

Crop Sequences 1 and 8

	(3) and (4)				(±2.38)
C1	31.4	36.3	41.0	38.7	36.9
C8	21.2	17.5	32.1	38.1	27.2
Mean	26.3	26.9	36.5	38.4	32.0
	(±1.56)				

Mean D.M. %: 79.9

(1) (±5.62) (3) (±3.05) For use in vertical and diagonal comparisons  
 (2) (±3.23) (4) (±2.20) For use in horizontal and interaction comparisons