Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



# Yields of the Field Experiments 1967



Full Table of Content

# 67/R/BQ/C/14 Previous Crops and N for Barley 1965-67 - Barley

# **Rothamsted Research**

Rothamsted Research (1968) 67/R/BQ/C/14 Previous Crops and N for Barley 1965-67 - Barley; Yields Of The Field Experiments 1967, pp 167 - 168 - **DOI**:

https://doi.org/10.23637/ERADOC-1-157

67/C/14.1

#### PREVIOUS CROPS X N FOR BARLEY

(BQ)

The effect of previous cropping and nitrogen on the yield of barley - Stackyard 1967, the third year - barley.

Design: A single replicate of 3 x 3 x 3 x 3 in 3 blocks of 9 plots, each split into 3 for N in 1966 and 1967.

Area of each sub plot: 0.0096. Area harvested: 0.0064.

Treatments: All combinations of:-

Whole plots (applied 1965):-

- 1. Cropping: Spring wheat (W), kale (K), Italian Ryegrass (G).
- Nitrogen: None (Q0), 1.0 (Q2), 2.0 (Q4) cwt N as 'Nitro-Chalk'.

Sub plots (applied to barley 1966 and 1967):-

- 3. 1966 Nitrogen: None (RO), 0.5 (R1), 1.0 (R2) N as 'Nitro-Chalk'.
- 4. 1967 Nitrogen: None (NO), 0.5 (N1), 1.0 (N2) N as 'Nitro-Chalk'.
- Basal applications: 2.5 cwt (0:20:20) combine drilled. Weedkiller: Mecoprop/2,4-D (Methoxone Extra at 6 pints in 35 gals).
- Cultivations, etc.: Ploughed twice: Sept 15, 1966 and Jan 3, 1967. Seed drilled at 165 lb: Mar 13. 'Nitro-Chalk' applied: Mar 13. Sprayed: May 11. Combine harvested: Aug 21. Variety: Maris Badger.
- NOTE: For previous years' results see 'Results' 65/C/27, 66/C/18.
- Standard errors per plot. Grain:
  Whole plot: 2.79 or 8.7% (16 d.f.)
  Sub plot: 3.44 or 10.8% (40 d.f.)

67/c/14.2

## SUMMARY OF RESULTS

### GRAIN

Crop in 1965	<b>Q</b> 0	N 1965 Q2	Q4	RO	RL.	R2	NO	Nl	N2	Mean
	(±1.61)			(1) and (2)			(1) and (2)			(±0.93)
W K G	31.2 30.1 33.2	28.8 30.7 35.5	32.1 29.2 36.9	30.4 30.0 35.2	29.9 28.8 35.2	31.8 31.1 35.1	14.7	33.1 32.4 40.1	41.5 42.8 45.1	30.0
				(1)	and (	2)	(1)	and (	2)	
		& Q2 Q4		32.4 30.6 32.7	30.1 32.1 31.6	31.9 32.3 33.9	17.7	34.7 34.5 36.5		31.7
				4 46.0		KGI		(±1.15	5)	(±0.66)
					RO R1 R2	St. S.	17.6 17.7 17.3	34.0 33.8 37.9	44.1 42.4 42.8	31.3
Mean (±0	.66)	4 (No.1)	agger 1			TABUTA	17.5	35.2	43.1	32.0

 <sup>(1) (±1.32)</sup> For use in vertical and diagonal comparisons
 (2) (±1.15) For use in horizontal and interaction comparisons

Mean D.M. %: 82.1