


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/R/EQ/C/16 Rate of Action of P Fertilisers 1965-67 - Ryegrass

Rothamsted Research

Rothamsted Research (1968) *67/R/EQ/C/16 Rate of Action of P Fertilisers 1965-67 - Ryegrass* ; Yields Of The Field Experiments 1967, pp 173 - 174 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/C/16.1

RYEGRASS AFTER BARLEY

(EQ)

The rate of action of P fertilisers, Sawyers II 1967, the third year.

Design: 4 replicates of 7 x 2 x 2 plus 2 plots per block without P, in 8 blocks of 16 plots.

Area of each plot: 0.0018. Area harvested: 0.0009.

Treatments 1965 and 1966: No P (0) (2 plots per block) and all combinations of:-

1. Phosphatic fertilisers:

Triple superphosphate	(S)
Potassium metaphosphate	(K)
Triple superphosphate plus potassium metaphosphate*	(SK)
Magnesium ammonium phosphate	(M)
Triple superphosphate plus magnesium ammonium phosphate*	(SM)
Potassium metaphosphate plus magnesium ammonium phosphate*	(KM)
Triple superphosphate plus potassium metaphosphate plus magnesium ammonium phosphate*	(SKM)

* Each material supplying half the P (one third in treatment SKM).

2. Levels of P: 12 lb (L1), 24 lb (L2) P to barley 1966 (double these rates to potatoes 1965).
3. Type of fertiliser: Powder, less than 1 mm (P), granular 1-4 mm (G).

All applied to barley 1966 with 'Nitro-Chalk', muriate of potash and kieserite in amounts adjusted to give a total of 60 lb N, 100 lb K and 25 lb Mg.

Basal applications: To ryegrass: 60 lb N and 50 lb K (16:0:16) for each cut.

Cultivations etc.: Basal NK applied: Feb 24, 1967. Cut once: May 17.

NOTE: For previous years' results see 'Results' 65/C/28 and 66/C/20.

Standard error per plot.

Dry matter: 2.84 or 5.8% (90 d.f.)

67/C/16.2

SUMMARY OF RESULTS

1ST AND ONLY CUT: DRY MATTER

	O	S	K	SK	M	SM	KM	SKM	Mean
Mean (±0.71)	45.8	48.8	51.6	49.9	48.8	49.4	50.3	50.4	49.4
	(±1.01)								(±0.38)
I1		49.2	50.0	47.6	48.5	49.5	50.2	46.7	48.8
I2		48.4	53.1	52.2	49.0	49.3	50.4	54.2	50.9
P		48.0	49.6	50.4	48.5	48.7	51.2	48.5	49.3
G		49.6	53.6	49.3	49.0	50.2	49.4	52.4	50.5

Mean D.M. %: 18.4