

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1967

[Full Table of Content](#)



### **67/R/ER/C/17 Rate of Action of P Fertilisers 1966-67 - Barley and Ryegrass**

#### **Rothamsted Research**

Rothamsted Research (1968) *67/R/ER/C/17 Rate of Action of P Fertilisers 1966-67 - Barley and Ryegrass* ; Yields Of The Field Experiments 1967, pp 175 - 176 - **DOI:**

**<https://doi.org/10.23637/ERADOC-1-157>**

67/c/17.1

BARLEY FOLLOWED BY RYEGRASS

(ER)

The rate of action of P fertilisers, Delharding, 1967, the second year.

Design: 3 replicates of 7 x 2 x 2 plus 2 plots per block without P, in 6 blocks of 16 plots.

Area of each plot:	Area harvested:
Barley: 0.0018	0.0009
Ryegrass: 0.0018	0.0006

Treatments 1966 and 1967: No P (0) (2 plots per block) and all combinations of:-

1. Phosphatic fertilisers:
  - Triple superphosphate (S)
  - Potassium metaphosphate (K)
  - Triple superphosphate plus potassium metaphosphate\* (SK)
  - Magnesium ammonium phosphate (M)
  - Triple superphosphate plus magnesium ammonium phosphate\* (SM)
  - Potassium metaphosphate plus magnesium ammonium phosphate\* (KM)
  - Triple superphosphate plus potassium metaphosphate plus magnesium ammonium phosphate\* (SKM)

\* Each material supplying half the P (one third in treatment SKM).

2. Levels of P: 12 lb (L1), 24 lb (L2) P to barley (double these rates to potatoes 1966).
3. Type of fertiliser: Powder, less than 1 mm (P), granular 1-4 mm (G).

All applied to barley with 'Nitro-Chalk', muriate of potash and kieserite in amounts adjusted to give a total of 60 lb N, 100 lb K and 25 lb Mg.

Basal applications: To barley: 60 lb N and 50 lb K (16:0:16).  
To ryegrass: 90 lb N as 'Nitro-Chalk'. Weedkillers: To barley: Dichlorprop/MCPA (Cornox RK Extra at 7 pints in 50 gals), paraquat to stubble at 1 lb ion in 40 gals.

Cultivations, etc.:

Barley: Ploughed: Sept 30, 1966. Fertilisers applied, seed drilled at 160 lb: Mar 15, 1967. Weedkiller applied: May 12. Basal NK applied: May 31. Harvested green: June 30. Paraquat applied: July 4. Variety: Maris Badger.

67/c/17.2

Ryegrass: Basal N applied, seed drilled at 60 lb: July 25, 1967.  
Cut: Oct 19. Variety: Italian S22.

NOTE: For previous years' results see 'Results' 66/c/21.

Standard errors per plot. Dry matter:  
Barley, green crop: 3.20 or 18.9% (60 d.f.)  
Ryegrass, 1st and only cut: 1.89 or 8.9% (60 d.f.)

SUMMARY OF RESULTS

	BARLEY GREENCROP. DRY MATTER. CWT PER ACRE								Mean
	O	S	K	SK	M	SM	KM	SKM	
Mean (±0.93)	6.3	17.7	17.8	20.1	16.7	19.7	17.1	20.2	17.0
				(±1.31)					(±0.43)
L1		16.8	14.8	17.3	13.1	16.7	13.6	16.1	15.5
L2		18.6	20.8	22.9	20.3	22.8	20.7	24.3	21.5
P		16.8	18.5	19.0	17.5	19.5	18.2	20.2	18.5
G		18.6	17.2	21.2	16.0	19.9	16.1	20.1	18.4

Mean D.M. %: 30.0

	RYEGRASS 1ST AND ONLY CUT. DRY MATTER.								Mean
	O	S	K	SK	M	SM	KM	SKM	
Mean (±0.54)	14.4	21.6	22.6	23.2	21.8	22.9	21.5	22.0	21.3
				(±0.77)					(±0.25)
L1		19.1	20.2	21.5	20.4	21.5	19.5	19.8	20.3
L2		24.2	25.0	25.0	23.1	24.3	23.5	24.1	24.2
P		20.6	22.1	23.4	20.6	23.4	21.8	22.1	22.0
G		22.7	23.0	23.0	23.0	22.4	21.3	21.9	22.5

Mean D.M. %: 11.4