


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/R/HWF/A/3 Hoosfield - Wheat After Fallow

Rothamsted Research

Rothamsted Research (1968) *67/R/HWF/A/3 Hoosfield - Wheat After Fallow* ; Yields Of The Field Experiments 1967, pp 19 - 19 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/A/3

WHEAT AFTER FALLOW - HOOSFIELD 1967

(HWF)

For history, treatments, etc. see 'Details' 1962.

The comparison of two varieties is now discontinued, Cappelle only being sown.

Area of each plot: 0.1238. Area harvested: 0.0368.

Cultivations, etc.:

Cropped plots: Ploughed: Sept 13, 1966. Seed drilled at 180 lb: Oct 28. Sprayed with ioxynil/mecoprop (Actril C at 6 pints in 20 gals): May 9, 1967. Combine harvested: Aug 30.

Fallow plots: Ploughed 3 times: Sept 13, 1966, Apr 3 and July 5, 1967.

SUMMARY OF RESULTS

Plot	B1	B2	B3
No of years of fallow	1	1	3
	GRAIN		
	15.9	18.1	22.2
	STRAW		
	11.9	12.1	13.4

Mean D.M. %: Grain: 85.2
Straw: 88.6