

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/S/SC/C/23 Intensive Wheat - Winter Wheat

Rothamsted Research

Rothamsted Research (1968) *67/S/SC/C/23 Intensive Wheat - Winter Wheat* ; Yields Of The Field Experiments 1967, pp 195 - 196 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/C/23.1

INTENSIVE WHEAT

(SC)

Saxmundham, Ollershaw's and Garner's plots 1967, the second year.

Area harvested: 0.0098.

The Meadow Fescue ley is now replaced by a one year Italian ryegrass ley, cut twice for hay, which receives 1.0 cwt N in the seedbed, and 0.5 cwt N and 1.0 cwt K₂O after the first cut. N as 'Nitro-Chalk', K₂O as muriate of potash.

Basal applications: 560 lb (0:20:20) broadcast before ploughing. Weedkiller to wheat: Mecoprop/2,4-D (Methoxone Extra at 7 pints in 20 gals). Insecticide to beans: Demeton-s-methyl at 3 oz in 37 gals.

Cultivations, etc.: Basal PK compound applied: Sept 20, 1966.

Ploughed: Sept 22.

Wheat: Seed drilled: Oct 17, 1966. 'Nitro-Chalk' applied: Mar 23, 1967. Weedkiller applied: Apr 27. Combine harvested: Aug 17. Variety: Cappelle.

Grass ley: Seed drilled: Mar 14, 1967. 'Nitro-Chalk' applied: Mar 23. Cut: June 13. 'Nitro-Chalk' and muriate of potash applied: June 26. Cut: Aug 8. Variety: S22 Italian.

Beans: Seed drilled at 270 lb: Mar 13, 1967. Insecticide applied: June 12. Combine harvested: Aug 24. Variety: Hasler's Throws.

NOTES: (1) Yields were taken for winter wheat only.

(2) Estimates of the incidence of take-all (*Ophiobolus graminis*) and eyespot (*Cercospora herpotrichoides*) were made.

Standard errors per plot. Grain:

Whole plot: 1.83 or 5.2% (6 d.f.)

Sub plot: 3.05 or 8.7% (18 d.f.)

67/C/23.2

SUMMARY OF RESULTS

Crop in 66 67	N1	N2	N3	Mean
GRAIN				
(1) and (2)				
				(±0.91)
W W	27.6	35.6	37.6	33.6
L W	36.2	43.7	39.4	39.8
W W	28.4	34.3	32.7	31.8
Mean (±0.88)	30.7	37.9	36.6	35.1

Mean D.M. %: 81.3

- (1) (±1.55) For use in vertical and diagonal comparisons
 (2) (±1.53) For use in horizontal and interaction comparisons

STRAW				
W W	22.1	30.5	30.4	27.7
L W	32.2	34.1	33.4	33.2
Mean	27.1	32.3	31.9	30.5

Mean D.M. %: 82.1