

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1967

[Full Table of Content](#)



---

### 67/R/AG/A/4 Agdell - Grass

#### Rothamsted Research

Rothamsted Research (1968) *67/R/AG/A/4 Agdell - Grass* ; Yields Of The Field Experiments 1967, pp 21 - 23 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/A/4.1

GRASS - AGDELL 1967

(AG)

For history, treatments etc. see 'Details' 1962 and 'Results' 63/A/4, 64/A/4, 65/A/4 and 66/A/4.

Area of each microplot: Plots 1-4 - 0.0180. Plots 5-6 - 0.0162.  
Area harvested: Plots 1-4 - 0.0023. Plots 5-6 - 0.0020.

P (as triple superphosphate) was applied in January 1967 to balance withdrawals by grass in 1964 and 1966 to all sub-plots except P0 which continued to receive no P.

K (as muriate of potash) was applied to balance withdrawals by grass in 1964, 1965 and 1966 to all sub-plots except K0 which continued to receive no K. Some of this K was applied in 1966 (see 'Results' 66/A/4). Because the amounts still required were large the 1967 dressings were split to lessen the risk of damaging the grass. Not more than 2.97 cwt K<sub>2</sub>O (3.32 cwt on sub-plots 5 and 6) - 10 lb muriate of potash per sub-plot - was applied in January and the balance was applied after the first cut but before ploughing.

Rates in cwt P<sub>2</sub>O<sub>5</sub>

Plot no.	Sub-plots testing P				Sub-plots testing K			
	P0	P1	P2	P4	K0	K1	K2	K4
1	0	0.75	0.85	0.93	0.80	1.00	0.88	0.91
2	0	0.68	0.75	0.83	0.59	0.86	0.83	0.88
3	0	0.71	0.70	0.89	0.70	0.76	0.84	0.73
4	0	0.57	0.60	0.80	0.73	0.87	0.83	0.86
5	0	0.52	0.60	0.72	0.50	0.72	0.76	0.80
6	0	0.50	0.69	0.76	0.51	0.80	0.70	0.80

Rates in cwt K<sub>2</sub>O

Plot no.	Sub-plots testing P				Sub-plots testing K			
	P0	P1	P2	P4	K0	K1	K2	K4
1	5.78	6.66	7.07	6.82	0	4.45	4.62	6.00
2	3.80	5.57	5.80	5.67	0	3.68	4.58	6.16
3	3.57	6.02	5.54	6.24	0	4.32	4.78	5.38
4	2.90	4.46	4.43	5.19	0	4.00	4.64	5.72
5	1.54	4.50	4.83	5.24	0	3.29	3.95	5.40
6	0.25	4.62	5.10	5.02	0	3.72	3.59	5.05

67/A/4.2

Basal dressing: 'Nitro-Chalk' applied at 0.8 cwt N in March.

Cultivations, etc.:-

Grass: P and first dressing of K applied: 18 Jan, 1967. Cut once for silage: 2 June, 1968. Second dressing of K applied: 7 June. Plots ploughed: 14 June. Rotary cultivated: 5 July, 21 July, 8 Aug, 24 Aug. Resown with Timothy: 7 Sept.  
Fallow: Ploughed: 31 Oct, 1966.

67/A/4.3

SUMMARY OF RESULTS

DRY MATTER

Plot

P	K	5	6	3	4	1	2	Mean
1ST AND ONLY CUT								
0	4	13.7	7.9	30.2	17.0	37.0	27.1	22.1
1	4	50.4	47.8	48.6	47.0	45.2	46.4	47.6
2	4	55.8	45.4	42.4	39.3	39.0	53.6	45.9
4	4	46.9	44.9	48.8	48.2	42.9	40.4	45.3
4	0	41.7	23.9	24.1	37.8	26.2	25.5	29.9
4	1	57.6	44.3	38.4	47.3	41.7	48.1	46.2
4	2	53.0	43.4	36.8	44.7	45.4	49.0	45.4
4	4	53.6	46.2	38.8	37.3	46.6	48.9	45.2
Mean		46.6	38.0	38.5	39.8	40.5	42.4	41.0

Mean D.M. %: 17.9