


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/R/CK/C/28 and 67/W/WCN/C/28 Potato Haulms - Potatoes

Rothamsted Research

Rothamsted Research (1968) *67/R/CK/C/28 and 67/W/WCN/C/28 Potato Haulms - Potatoes* ; Yields Of The Field Experiments 1967, pp 211 - 214 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/c/28.1

EFFECT OF POTATO HAULM ON WINTER WHEAT

(CK and WCN)

Rothamsted (R) Pastures and Woburn (W) Broadmead II, 1967, the first year (potatoes).

Design: 4 blocks of 8 plots, randomisation restricted.

Area of each plot: 0.0161. Area harvested: Pastures (R) - 0.0096, Broadmead II (W) - 0.0099.

Treatments: All combinations of:-

1. Nitrogen to potatoes: None (NO), 0.66 (N2), 1.32 (N4), 2.0 (N6) as 'Nitro-Chalk'.
2. Haulm disposal: Burnt off with acid (B), cut and removed (C).

Basal applications: 1200 lb (0:14:28) broadcast by drill.

Pastures (R): Weedkiller: Linuron at 0.75 lb plus paraquat at 0.5 lb ion in 37 gals. Fungicide: Mancozeb at 1.2 lb in 30 gals on four occasions.

Broadmead II (W): Weedkiller: Linuron at 0.5 lb plus paraquat at 0.5 lb ion in 50 gals. Insecticide: Ground application - demeton-s-methyl 3 oz in 30 gals, aerial application (by fixed wing aircraft) 3 oz in 2 gals. Fungicide: Mancozeb at 1.2 lb in 30 gals on 3 occasions.

Cultivations, etc.:-

Pastures (R): Rotary cultivated: Sept 16, 1966. Ploughed: Nov 23 - Jan 5, 1967. Basal PK applied: Mar 31. 'Nitro-Chalk' applied: Apr 4. Rotary cultivated second time, potatoes planted: Apr 6. Weedkiller applied: May 12. Fungicide applied: June 29, July 21, Aug 3, Aug 8. Haulm removed from C plots: Aug 29. B plots sprayed with undiluted BOV at 15 gals: Sept 7. Haulm destroyed mechanically: Sept 14. Lifted: Sept 15. Variety: Pentland Dell. Previous crops: Barley 1965, fallow 1966.

Broadmead II (W): Ploughed: Oct 22, 1966. Ground chalk applied at 4.5 tons: Jan 9, 1967. Basal PK applied: Mar 20. 'Nitro-Chalk' applied: Mar 28. Potatoes planted: Mar 30. Weedkiller applied: May 3. Rotary ridged: June 9. Insecticide and fungicide applied: June 20. Fungicide applied: June 30, Aug 8.

67/C/28.2

Insecticide applied from the air: July 13. Haulm removed from C plots: Aug 24. B plots sprayed with diquat (Reglone at 4 pints in 40 gals): Aug 25. Lifted: Sept 11. Variety: Pentland Dell. Previous crops: Winter wheat 1965, spring wheat 1966.

NOTE: Samples of haulm were taken from C plots at the time of cutting for dry matter and percentage N. Potatoes were sampled at harvest for dry matter and percentage N.

Standard errors per plot. Total tubers:
Pastures (R): 0.925 or 5.0% (21 d.f.)
Broadmead II (W): 1.198 or 11.7% (21 d.f.)

67/c/28.3

SUMMARY OF RESULTS

	TOTAL TUBERS				Mean
	N0	N2	N4	N6	
	PASTURES (R)				
	(±0.463)				(±0.231)
B	15.91	17.96	20.10	20.93	18.73
C	14.68	19.25	19.10	19.53	18.14
Mean (±0.327)	15.29	18.60	19.60	20.23	18.43
	BROADMEAD II (W)				
	(±0.599)				(±0.299)
B	7.85	9.24	12.96	12.06	10.53
C	6.43	9.98	10.95	12.33	9.92
Mean (±0.423)	7.14	9.61	11.96	12.19	10.23

67/c/28.4

		% WARE				
		NO	N2	N4	N6	Mean
PASTURES (R)						
B		98.2	98.5	98.1	98.1	98.2
C		97.6	98.2	98.7	98.1	98.1
Mean		97.9	98.3	98.4	98.1	98.2
BROADMEAD II (W)						
B		90.6	94.9	97.1	96.9	94.9
C		86.9	94.6	96.4	97.3	93.8
Mean		88.8	94.8	96.8	97.1	94.3