

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/R/EAT/C/33 N and 67/W/WEAQ/C/33 N - Fixation - Lucerne and Grass

Rothamsted Research

Rothamsted Research (1968) *67/R/EAT/C/33 N and 67/W/WEAQ/C/33 N - Fixation - Lucerne and Grass* ; Yields Of The Field Experiments 1967, pp 227 - 236 - DOI:

<https://doi.org/10.23637/ERADOC-1-157>

67/C/33.1

N FIXATION - LUCERNE AND GRASS

(EAT and WEAQ)

Nitrogen fixation by lucerne - Rothamsted (R) Stackyard and Woburn (W) Stackyard C 1967, the first year.

Design: 4 blocks of four plots split into four for minerals and nitrogen.

Area of each sub-plot: 0.0011. Area harvested: 0.0006.

Treatments: All combinations of:-

- Whole plots: 1. Seed: Lucerne uninoculated (O), lucerne with ineffective inoculum (I), lucerne with effective inoculum 2001 (E), Italian ryegrass (G).
- Sub-plots: 2. Minerals: Unmanured (U), 0.6 cwt P₂O₅, 1.2 cwt K₂O as compound fertiliser (0:14:28), plus 4 tons of ground chalk (5 tons on Stackyard C (W)) (M).
3. Nitrogen (as 'Nitro-Chalk'): None (NO), 60 lb N* before the first cut (N1), 60 lb N* before the first cut, 60 lb after the first cut (N2), 60 lb N* before the first cut, 60 lb after the first cut, 60 lb after the second cut (N3).

*30 lb in the seedbed, 30 lb in June.

Basal applications: None.

Cultivations, etc.:-

Stackyard (R): Ploughed: Nov 16, 1966. PK, ground chalk and seedbed 'Nitro-Chalk' applied, plots rotary cultivated, seed drilled, lucerne at 12 lb, ryegrass at 30 lb: Apr 26, 1967. 'Nitro-Chalk' applied: June 27. Cut three times: June 24, Aug 29, Oct 25. Appropriate 'Nitro-Chalk' dressings applied after first two cuts. Varieties: Lucerne - Du Puits, ryegrass S22. Previous crops: Barley 1965, barley 1966.

Stackyard C (W): Ploughed: Oct 28, 1966. PK, ground chalk and seedbed 'Nitro-Chalk' applied, plots rotary cultivated, seed drilled, lucerne at 12 lb, ryegrass at 30 lb: Apr 27. 'Nitro-Chalk' applied: June 28. Cut three times: July 26, Sept 4, Oct 24. Appropriate 'Nitro-Chalk' dressings applied after first two cuts. Varieties: Lucerne - Du Puits, ryegrass S22. Previous crops: Fallow 1965 and 1966.

NOTE: Soil samples were taken to determine pH before sowing. Rhizobium counts were taken from root samples on each experiment.

67/C/33.2

Standard errors per plot. Dry matter:

Stackyard (R): 1st cut:	Whole plot: 3.58 or 25.0% (6 d.f.)
	Sub plot: 2.20 or 15.4% (24 d.f.)
2nd cut:	Whole plot: 2.06 or 10.9% (6 d.f.)
	Sub plot: 2.77 or 14.6% (24 d.f.)
3rd cut:	Whole plot: 0.42 or 7.8% (6 d.f.)
	Sub plot: 1.91 or 35.3% (24 d.f.)
Total of 3 cuts:	Whole plot: 4.81 or 12.5% (6 d.f.)
	Sub plot: 4.87 or 12.6% (24 d.f.)

Stackyard C (W): 1st cut:	Whole plot: 5.36 or 32.4% (6 d.f.)
	Sub plot: 4.38 or 26.5% (24 d.f.)
2nd cut:	Whole plot: 2.74 or 20.7% (6 d.f.)
	Sub plot: 2.68 or 20.2% (24 d.f.)
3rd cut:	Whole plot: 3.15 or 43.5% (6 d.f.)
	Sub plot: 1.60 or 22.1% (24 d.f.)
Total of 3 cuts:	Whole plot: 9.01 or 24.3% (6 d.f.)
	Sub plot: 7.65 or 20.7% (24 d.f.)

67/C/33.3

SUMMARY OF RESULTS

DRY MATTER

	O	I	E	G	Mean
STACKYARD (R)					
1ST CUT					
Mean (± 1.79)	8.8	5.6	20.4	22.4	14.3
		(1) and (2)			(± 0.39)
U	7.6	5.1	17.4	22.0	13.0
M	10.1	6.1	23.4	22.8	15.6
		(3) and (4)			(± 0.55)
NO	7.0	3.6	19.7	11.4	10.4
N1	9.1	7.1	20.0	26.2	15.6
N2	8.8	6.6	21.7	24.5	15.4
N3	10.3	5.1	20.1	27.5	15.8

Mean D.M. %: 27.8

(1) (± 1.87) (3) (± 2.03) For use in horizontal and diagonal comparisons only
 (2) (± 0.78) (4) (± 1.10) For use in vertical and interaction comparisons only

NOTE: 1st cut N1 = N2 = N3

67/c/33.4

DRY MATTER					
	O	I	E	G	Mean
STACKYARD (R)					
2ND CUT					
Mean (± 1.03)	20.3	16.0	24.3	15.0	18.9
	(1) and (2)				(± 0.49)
U	18.4	14.4	21.7	15.7	17.6
M	22.2	17.5	26.8	14.3	20.2
	(3) and (4)				(± 0.69)
N0	15.4	6.9	23.0	4.5	12.5
N1	18.7	14.2	23.2	7.9	16.0
N2	23.6	21.9	25.2	24.7	23.9
N3	23.4	20.8	25.7	22.8	23.2

Mean D.M. %: 19.7

(1) (± 1.24) (3) (± 1.58) For use in horizontal and diagonal comparisons only
 (2) (± 0.98) (4) (± 1.38) For use in vertical and interaction comparisons only

NOTE: 2nd cut N2 = N3

67/C/33.5

DRY MATTER					
	O	I	E	G	Mean
STACKYARD (R)					
3RD CUT					
Mean (± 0.21)	6.6	4.5	6.6	4.0	5.4
		(1) and (2)			(± 0.34)
U	5.5	4.1	5.7	4.3	4.9
M	7.7	4.9	7.5	3.7	5.9
		(3) and (4)			(± 0.48)
NO	6.2	3.8	7.7	1.6	4.8
N1	5.1	3.6	6.3	1.7	4.2
N2	6.7	3.5	5.9	3.2	4.8
N3	8.3	7.0	6.6	9.4	7.8

Mean D.M. %: 20.9

(1) (± 0.52) (3) (± 0.85) For use in horizontal and diagonal comparisons only
 (2) (± 0.68) (4) (± 0.95) For use in vertical and interaction comparisons only

67/c/33.6

DRY MATTER					
	O	I	E	G	Mean
STACKYARD (R)					
TOTAL OF 3 CUTS					
Mean (± 2.40)	35.7	26.0	51.3	41.4	38.6
	(1) and (2)				(± 0.86)
U	31.4	23.6	44.8	42.0	35.4
M	40.0	28.4	57.8	40.8	41.7
	(3) and (4)				(± 1.22)
NO	28.7	14.3	50.4	17.5	27.7
N1	32.9	24.9	49.5	35.8	35.8
N2	39.1	32.0	52.7	52.4	44.1
N3	42.1	32.9	52.4	59.8	46.8

Mean D.M. %: 22.8

(1) (± 2.69) (3) (± 3.20) For use in horizontal and diagonal comparisons only
 (2) (± 1.72) (4) (± 2.43) For use in vertical and interaction comparisons only

67/C/33.7

DRY MATTER					
	O	I	E	G	Mean
STACKYARD C (W)					
1ST CUT					
Mean (± 1.34)	14.0	13.1	19.0	20.0	16.5
		(1) and (2)			(± 0.77)
U	11.9	13.1	16.5	19.9	15.3
M	16.2	13.1	21.5	20.2	17.8
		(3) and (4)			(± 1.09)
N0	12.4	8.4	16.9	7.8	11.4
N1	15.8	13.8	21.6	23.4	18.7
N2	12.1	14.7	20.3	25.7	18.2
N3	15.8	15.4	17.3	23.3	17.9

Mean D.M. %: 26.6

(1) (± 1.73) (3) (± 2.32) For use in horizontal and diagonal comparisons only
 (2) (± 1.55) (4) (± 2.19) For use in vertical and interaction comparisons only

NOTE: 1st cut N1 = N2 = N3

67/c/33.8					
DRY MATTER					
	O	I	E	G	Mean
STACKYARD C (W)					
2ND CUT					
Mean (± 0.68)	14.0	12.7	16.6	9.7	13.2
	(1) and (2)				(± 0.47)
U	12.1	11.7	14.6	9.3	11.9
M	15.9	13.7	18.7	10.1	14.6
	(3) and (4)				(± 0.67)
N0	12.4	9.2	13.9	2.6	9.5
N1	14.6	10.2	17.2	4.5	11.7
N2	14.4	16.2	18.3	15.2	16.0
N3	14.5	15.2	17.1	16.3	15.8

Mean D.M. %: 26.2

(1) (± 0.96) (3) (± 1.35) For use in horizontal and diagonal comparisons only
 (2) (± 0.95) (4) (± 1.34) For use in vertical and interaction comparisons only

NOTE: 2nd cut N2 = N3

67/c/33.9

DRY MATTER					
	O	I	E	G	Mean
STACKYARD C (W)					
3RD CUT					
Mean (± 0.79)	8.3	6.5	8.7	5.5	7.2
		(1) and (2)			(± 0.28)
U	7.2	5.4	8.3	5.3	6.5
M	9.4	7.5	9.2	5.7	7.9
		(3) and (4)			(± 0.40)
NO	6.8	5.4	7.5	1.2	5.2
N1	8.8	4.9	9.0	1.7	6.1
N2	7.7	7.2	9.2	3.7	6.9
N3	9.9	8.4	9.2	15.3	10.7

Mean D.M. %: 19.9

(1) (± 0.88) (3) (± 1.05) For use in horizontal and diagonal comparisons only
 (2) (± 0.57) (4) (± 0.80) For use in vertical and interaction comparisons only

67/c/33.10

DRY MATTER					
	O	I	E	G	Mean
STACKYARD C (W)					
TOTAL OF 3 CUTS					
Mean (± 2.25)	36.3	32.2	44.4	35.2	37.0
		(1) and (2)			(± 1.35)
U	31.2	30.2	39.3	34.4	33.8
M	41.5	34.3	49.4	35.9	40.3
		(3) and (4)			(± 1.91)
N0	31.7	23.0	38.3	11.5	26.1
N1	39.3	29.0	47.8	29.6	36.4
N2	34.2	38.1	47.8	44.6	41.2
N3	40.1	38.9	43.5	54.9	44.4

Mean D.M. %: 24.2

(1) (± 2.95) (3) (± 4.01) For use in horizontal and diagonal comparisons only
 (2) (± 2.71) (4) (± 3.83) For use in vertical and interaction comparisons only