


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/W/WEAM/C/34 Placement of Fumigant

Rothamsted Research

Rothamsted Research (1968) *67/W/WEAM/C/34 Placement of Fumigant* ; Yields Of The Field Experiments 1967, pp 237 - 238 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/C/34.1

PLACEMENT OF FUMIGANT FOR POTATOES

(WEAM)

Economic row and planting-station fumigation for control of potato cyst-nematode - Woburn Butt Furlong 1967.

Design: 3 randomised blocks of 5 plots.

Area of each plot: 0.0024. Area harvested: 0.0024.

Treatments:

Fumigants: DD: None (D0), 56 (D1), 112 (D2), 224 lb (D4) placed in the row by gravity feed, and 140 lb (DH) injected into the row at 18 in. intervals.

Basal applications: 10 cwt (17:11:22). Weedkiller: Paraquat at 0.5 lb plus linuron at 0.25 lb in 30 gals. Fungicide: Mancozeb 1.2 lb in 37 gals on 2 occasions.

Cultivations, etc.: Ploughed: Dec 19, 1966. Basal NPK applied: Mar 3, 1967. Rotary cultivated, rotary ridged, DD applied in the ridge: Mar 7. Potatoes hand planted in ridges at 18 inch intervals: Apr 19. Sprayed weedkiller: May 9. Rotary ridged: June 9. Sprayed fungicide: July 1 and July 26. Lifted: Sept 6. Variety: Majestic. Previous crops: Barley 1965, potatoes 1966.

Standard error per plot.

Total tubers: 0.668 or 8.9% (8 d.f.)

67/C/34.2

SUMMARY OF RESULTS

DO	D1	D2	D4	DH	Mean
TOTAL TUBERS					
2.21	6.62	8.54 (±0.386)	12.29	7.92	7.52
% WARE					
51.9	68.7	76.4	81.0	71.3	69.9