

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1967

[Full Table of Content](#)



---

### 67/W/WECG/C/41 Cereal Cyst Nematode - Barley and Spring Wheat

#### Rothamsted Research

Rothamsted Research (1968) *67/W/WECG/C/41 Cereal Cyst Nematode - Barley and Spring Wheat ; Yields Of The Field Experiments 1967*, pp 259 - 260 - DOI:

<https://doi.org/10.23637/ERADOC-1-157>

67/c/41.1

CEREAL CYST - NEMATODE

(WECG)

The effect of cereal cyst-nematode (*Heterodera avenae*) on the yield of spring wheat and barley - Butt Close 1967.

Design: (Each crop) 3 blocks of 5 split into 3. Blocks arranged in pairs, one of each crop.

Area of each sub plot: 0.0032. Area harvested: 0.0021.

Treatments: All combinations of:-

Whole plots: 1. Oats 1966: Under oats till harvest - (C,D,E) (3 plots per block). Oats rotary cultivated May 26 and then bare fallowed (A). Oats rotary cultivated May 26, bare fallowed, injected with D-D at 400 lb Dec 19, 1966 (B).

Sub plots: 2. Nitrogen to wheat 1967: 0.6 (N1), 1.2 (N2), 1.8 (N3) cwt N as 'Nitro-Chalk'.  
3. Nitrogen to barley 1967: 0.4 (N1), 0.8 (N2), 1.2 (N3) cwt N as 'Nitro-Chalk'.

Basal application: Oats 1966: 2 cwt (21:14:14) combine drilled. Weedkiller. Ioxynil/mecoprop (Actril C at 5 pints in 35 gals). Wheat and barley 1967: 280 lb (0:20:20). Weedkiller: Amino-triazole at 4 lb plus ammonium thiocyanate at 3.7 lb in 33 gals.

Cultivations, etc.:-

Oats 1966: Ploughed: 15 - 23 Nov, 1965. Seed combine drilled at 170 lb: Mar 9, 1966. Weedkiller applied: May 13. A and B plots rotary cultivated: May 26, Aug 18. Combine harvested: Sept 12. Variety: Condor. Previous crops: Barley 1964, carrots, kale, cabbage, 1965.

Wheat and barley 1967: Weedkiller applied: Sept 27, 1966. Ground chalk applied at 20 cwt: Oct 31. Ploughed: Nov 2. DD applied to B plots: Dec 19. Seed combine drilled, wheat at 170 lb, barley at 140 lb: Mar 6, 1967. 'Nitro-Chalk' applied: Mar 13. Combine harvested: Aug 23. Varieties: Wheat - Kloka, barley - Maris Badger.

NOTE: Soil samples were taken for nematode counts on June 5.

Standard errors per plot. Grain:

Spring wheat.	Whole plot: 2.64 or 7.2% (8 d.f.)
	Sub plot: 4.58 or 12.6% (20 d.f.)
Barley.	Whole plot: 2.22 or 6.3% (8 d.f.)
	Sub plot: 3.22 or 9.1% (20 d.f.)

67/C/41.2

SUMMARY OF RESULTS

	GRAIN					Mean
	A	B	C	D	E	
	SPRING WHEAT					
	(1) and (2)					(±1.18)
N1	35.3	42.2	32.7	34.5	31.4	35.2
N2	37.2	42.1	42.0	34.2	33.6	37.8
N3	38.1	38.9	35.2	34.5	33.9	36.1
Mean (±1.52)	36.9	41.1	36.6	34.4	33.0	36.4

Mean D.M. %: 83.8

(1) (±2.64) For use in horizontal and diagonal comparisons

(2) (±2.64) For use in vertical and interaction comparisons

	BARLEY					Mean
	A	B	C	D	E	
	(1) and (2)					(±0.83)
N1	33.5	36.3	28.3	24.8	25.7	29.7
N2	38.6	34.5	38.5	39.7	36.3	37.5
N3	37.4	41.0	37.5	37.4	38.7	38.4
Mean (±1.28)	36.5	37.3	34.8	33.9	33.6	35.2

Mean D.M. %: 83.5

(1) (±1.99) For use in horizontal and diagonal comparisons

(2) (±1.86) For use in vertical and interaction comparisons